

Finite Volume and Partially Quenched QCD-like Effective Field Theories

Johan Bijnens and Thomas Rössler

Department of Astronomy and Theoretical Physics, Lund University,
Sölvegatan 14A, SE 223-62 Lund, Sweden

Abstract

We present a calculation of the meson masses, decay constants and quark-antiquark vacuum expectation value for the three generic QCD-like chiral symmetry breaking patterns $SU(N_F) \times SU(N_F) \rightarrow SU(N_F)_V$, $SU(N_F) \rightarrow SO(N_F)$ and $SU(2N_F) \rightarrow Sp(2N_F)$ in the effective field theory for these cases. We extend the previous two-loop work to include effects of partial quenching and finite volume.

The calculation has been performed using the quark flow technique. We reproduce the known infinite volume results in the unquenched case. The analytical results can be found in the supplementary material.

Some examples of numerical results are given. The numerical programs for all cases are included in version 0.54 of the CHIRON package.

The purpose of this work is the use in lattice extrapolations to zero mass for QCD-like and strongly interacting Higgs sector lattice calculations.

1 Introduction

Effective field theory is used extensively in the study of strongly interacting gauge theories. A recent review covering a number of different applications in addition to other methods is [1]. Besides general interest in understanding strongly interacting gauge theories, they might still be useful as an alternative for the Standard Model Higgs sector as well as for dark matter. These applications have been reviewed recently at the 2015 [2, 3] and 2013 [4] lattice conferences. A number of recent lattice studies is [5]. Reviews of technicolor and strongly interacting Higgs sectors are [6, 7, 8].

Lattice studies are always performed at a nonzero fermion mass. In order to obtain results in the massless limit extrapolations are needed. A main tool for this in the context of lattice QCD is Chiral Perturbation Theory (ChPT) [9, 10, 11].

In the case of equal mass fermions three main symmetry breaking patterns are possible [12, 13, 14]. For N_F Dirac fermions in a complex representation the global symmetry group is $SU(N_F)_L \times SU(N_F)_R$ and it breaks spontaneously to the diagonal subgroup $SU(N_F)_V$. For N_F Dirac fermions in a real representation the global symmetry group is $SU(2N_F)$ and it breaks spontaneously to $SO(2N_F)$. An alternative possibility is that we have N_F Majorana fermions in a real representation with a global symmetry group $SU(N_F)$ spontaneously broken to $SO(N_F)$. We show in this work that the EFT for the quantities we consider is really the same as for Dirac fermions. The final case is N_F Dirac fermions in a pseudo-real representation. The global symmetry group is again $SU(2N_F)$ but in this case it is expected to be broken spontaneously to $Sp(2N_F)$.

The effective field theory (EFT) for these cases is discussed at tree level or lowest order (LO) in [15]. At next-to-leading order (NLO) the first case is simply ChPT for N_F light quarks with a symmetry breaking pattern of $SU(N_F) \times SU(N_F) \rightarrow SU(N_F)$, a direct extension of the QCD case and was already done in [11]. The pseudo-real case was done at NLO by [16]. The $SU(2N_F) \rightarrow SO(2N_F)$ case was done in [17]. The extension for all three cases to next-to-next-to-leading order (NNLO) was done in earlier work by one of the authors [18]. More references to earlier work can be found there and in [19, 20].

This paper is an extension to the work of [18]. We add a short discussion showing that the calculations and the Lagrangian for the real case also covers the case of Majorana fermions in a real representation. The main part of the work concerns the extension of the calculations at NNLO order of the masses, decay constants and vacuum expectation values to include effects of partial quenching and finite volume.

Partial quenching was introduced in ChPT by [21]. A thorough discussion of the assumptions involved is in [22]. It allows to study a number of variations of input parameters at reduced cost, as discussed in e.g. [23]. We do not use the supersymmetric method introduced in [21] and extended (at NLO) to the cases discussed here in [17]. We only use the quark-flow technique introduced in [24]. Two-loop results in infinite volume partially quenched ChPT (PQChPT) for the masses and decay constants are in [25, 26, 27]. The definitions of the infinite volume integrals we use can be found there.

Finite volume effects in ChPT were introduced in ChPT in [28, 29, 30]. Early two-loop work is [31, 32]. The vacuum expectation value was discussed in more detail in [33].

After the proper evaluation of the finite volume two-loop sunset integrals using two different methods [34] the masses and decay constants were treated in both the unquenched [35] and partially quenched [36] case. In particular the integral notation at finite volume we use is defined in [36].

In Sect. 2 we recapitulate briefly the discussion from [18] at the quark level and add the case with Majorana fermions. Sect. 3 similarly recapitulates [18] at the effective field theory level and adds the Majorana fermion case. The cases with Dirac fermions and Majorana fermions are essentially identical from the EFT point of view for the quantities we consider. The underlying reason is an $U(2N_F)$ transformation that relates the two cases as discussed in Sect. 4. Partial quenching and the quark.flow techniques we have used for the different cases is discussed to some extent in Sect. 5. For a discussion on finite volume and the notation used there we refer to [36]. Our analytical results are described in Sect. 6, in particular we clarify the definitions of the decay constant and vacuum expectation value used in terms of quark fields. The numerical examples and checks are presented in Sect. 7. The analytical formulas are included in the supplementary file [37] and the numerical programs are available via CHIRON, [38, 39]. The last section briefly recapitulates the main points of our work.

2 Quark level

2.1 The three Dirac fermion cases

The discussion here is kept very short, longer versions can be found in [15] and [18]. This subsection is mainly included to show normalization conventions.

QCD or complex representation In the N_F equal mass Dirac fermions in a complex representation, we put the N_F fermions together in an N_F column matrix q . The global symmetry transformation by $g_L \times g_R \in SU(N_F)_L \times SU(N_F)_R$ is given by

$$q_L \rightarrow g_L q_L, \quad q_R \rightarrow g_R q_R, \quad l_\mu \rightarrow g_L l_\mu g_L^\dagger + i g_L \partial_\mu g_L^\dagger, \quad r_\mu \rightarrow g_R l_\mu g_R^\dagger + i g_R \partial_\mu g_R^\dagger, \quad \mathcal{M} \rightarrow g_R \mathcal{M} g_L^\dagger. \quad (1)$$

The matrix $\mathcal{M} = m_q \mathbf{I} + s + ip$ brings the quark mass term $m_q \mathbf{I}$ and the external scalar s and pseudo-scalar densities in the Lagrangian via $-\bar{q}_R \mathcal{M} q_L + \text{h.c.}$. The external fields l_μ, r_μ are in the Lagrangian via $\bar{q}_L \gamma^\mu l_\mu q_L + \bar{q}_R \gamma^\mu r_\mu q_R$. Taking derivatives w.r.t. the external fields allows to calculate relevant Green functions [10, 11]. In particular, deriving w.r.t. s_{11} allows us to obtain $\langle \bar{q}_{L1} q_{R1} + \bar{q}_{R1} q_{L1} \rangle$ and derivatives w.r.t. $a_{\mu 12}$ with $r_{\mu 12} = -l_{\mu 12} = a_{\mu 12}$ allows access to matrix-elements of $\bar{q}_2 \gamma^\mu \gamma^5 q_1$. The symmetry is spontaneously broken by a vacuum expectation value

$$\langle \bar{q}_{Lj} q_{Ri} \rangle = v_0 \delta_{ij}. \quad (2)$$

This leaves a global symmetry $SU(N_F)_V$ with $g_L = g_R$ unbroken.

Adjoint or real representation When the fermions are in a real representation, we can introduce besides the N_F right handed fermions q_{Ri} a second set of right handed fermions in the same gauge group representation, $\tilde{q}_{Ri} = C\bar{q}_{Li}^T$. These can be put together in a $2N_F$ column vector \hat{q} , $\hat{q}^T = (q_{R1} \dots q_{RN_F} \tilde{q}_{R1} \dots \tilde{q}_{RN_F})$. The global symmetry transformation with $g \in SU(2N_F)$ is now

$$\hat{q} \rightarrow g\hat{q}, \quad \hat{V}_\mu \rightarrow g\hat{V}_\mu g^\dagger, \quad \hat{\mathcal{M}} \rightarrow g\hat{\mathcal{M}}g^T. \quad (3)$$

We define the external densities and currents as in the QCD case with r_μ, l_μ and \mathcal{M} . We define $2N_F \times 2N_F$ matrices

$$\hat{\mathcal{M}} = \begin{pmatrix} 0 & \mathcal{M} \\ \mathcal{M}^T & 0 \end{pmatrix}, \quad \hat{V}_\mu = \begin{pmatrix} r_\mu & 0 \\ 0 & -l_\mu^T \end{pmatrix}. \quad (4)$$

Note that the global symmetry can change quark-antiquark currents to diquark currents. The fermions condense forming a vacuum expectation value

$$\frac{1}{2} \langle \hat{q}_j^T C \hat{q}_j \rangle = v_0 J_{Sij} \quad J_S = \begin{pmatrix} 0 & \mathbf{I} \\ \mathbf{I} & 0 \end{pmatrix}. \quad (5)$$

This leaves a global symmetry $SO(2N_F)$ with $gJ_S g^T = \mathbf{I}$.

$N_c = 2$ or pseudo-real representation When the fermions are in a pseudo-real representation, we can introduce besides the N_F right handed fermions q_{Ria} again a second set of right handed fermions in the same gauge group representation, $\tilde{q}_{Ria} = \epsilon_{ab} C\bar{q}_{Lib}^T$. a, b are gauge indices and the extra Levi-Civita tensor ϵ_{ab} is needed to have \tilde{q}_{Ria} transform under the gauge group as q_{iRa} . The explicit formula is for the case of the fundamental representation with $N_c = 2$. q_{Ri} and \tilde{q}_{Ri} can be put together in a $2N_F$ column vector \hat{q} , $\hat{q}^T = (q_{R1} \dots q_{RN_F} \tilde{q}_{R1} \dots \tilde{q}_{RN_F})$. The global symmetry transformation with $g \in SU(2N_F)$ is now

$$\hat{q} \rightarrow g\hat{q}, \quad \hat{V}_\mu \rightarrow g\hat{V}_\mu g^\dagger, \quad \hat{\mathcal{M}} \rightarrow g\hat{\mathcal{M}}g^T. \quad (6)$$

We define the external densities and currents as in the QCD case with r_μ, l_μ and \mathcal{M} . We then define

$$\hat{\mathcal{M}} = \begin{pmatrix} 0 & -\mathcal{M} \\ \mathcal{M}^T & 0 \end{pmatrix}, \quad \hat{V}_\mu = \begin{pmatrix} r_\mu & 0 \\ 0 & -l_\mu^T \end{pmatrix}. \quad (7)$$

Note that the global symmetry can again change quark-antiquark currents to diquark currents. The fermions condense forming a vacuum expectation value

$$\frac{1}{2} \langle \hat{q}_{ja}^T \epsilon_{ab} C \hat{q}_{jb} \rangle = v_0 J_{Aij} \quad J_A = \begin{pmatrix} 0 & -\mathbf{I} \\ \mathbf{I} & 0 \end{pmatrix}. \quad (8)$$

This leaves a global symmetry $Sp(2N_F)$ with $gJ_A g^T = \mathbf{I}$.

2.2 Majorana fermions in a real representation

In the earlier work [18] at infinite volume Dirac fermions and Dirac masses were assumed. It was then also assumed that the vacuum condensate was aligned with the Dirac fermion masses. There is in fact another possibility. Majorana fermions with a Majorana mass in a real representation of the gauge group. In this case the global symmetry is $SU(N_F)$. It is expected to be spontaneously broken down to $SO(N_F)$ which is aligned with the Majorana masses.

A Majorana spinor is a Dirac spinor that satisfies

$$\psi = C\bar{\psi}^T \quad \text{or} \quad \psi = \begin{pmatrix} \psi_M \\ -i\sigma^2\psi_M^* \end{pmatrix}. \quad (9)$$

The last equality are in the chiral representation for the Dirac matrices. The Lagrangian for a single free Majorana fermion is

$$\frac{1}{2}\bar{\psi}i\gamma^\mu\partial_\mu\psi - \frac{m}{2}\bar{\psi}\psi = \psi_M^\dagger Ci\bar{\sigma}^\mu\partial_\mu\psi - \frac{im}{2}(\psi_M^T\sigma^2\psi + \psi_M^\dagger\sigma^2\psi^*). \quad (10)$$

$\bar{\sigma}^0 = \text{I}$, $\bar{\sigma}^i = -\sigma^i$. If we want to gauge this for $m \neq 0$ the mass term requires the fermions to be in a real representation of the gauge group.

For N_F Majorana fermions ψ_{Mi} in the adjoint representation with external fields \hat{V}_μ and $\hat{\mathcal{M}}$ the Lagrangian, put in a big column vector $\hat{q}^T = (\psi_1^T \dots \psi_{N_F}^T)$ is

$$\mathcal{L} = \frac{1}{2}\text{tr}_c(\hat{q}^\dagger i\bar{\sigma}^\mu(iD_\mu + \hat{V}_\mu)\hat{q}) - \frac{1}{2}\text{tr}_c(\hat{q}^T\sigma^2\hat{\mathcal{M}}^\dagger\hat{q} + \hat{q}^\dagger\sigma^2\hat{\mathcal{M}}\hat{q}^*). \quad (11)$$

This Lagrangian has a global $SU(N_F)$ symmetry with $g \in SU(N_F)$ with

$$\hat{q} \rightarrow g\hat{q}, \quad \hat{V}_\mu \rightarrow g\hat{V}_\mu g^\dagger + ig\partial_\mu g^\dagger, \quad \hat{\mathcal{M}} \rightarrow g\hat{\mathcal{M}}g^T. \quad (12)$$

The maximal symmetry argument says that in this case the fermions will condense to the flavour neutral vacuum $\langle \text{tr}_c(\hat{q}^T C \hat{q}) \rangle$. This is conserved by the part of the global group that satisfies $gg^T = \text{I}$ or the conserved part of the global symmetry group is $SO(N_F)$.

Note that the form of the vacuum and the form of the mass term are the only differences as far as the global symmetry group and its breaking are concerned compared to the case with $N_F/2$ Dirac fermions in a real representation.

3 Effective field theory

3.1 The general LO and NLO Lagrangian

The ChPT Lagrangian for N_F flavours at LO and NLO has been derived in [11]. The Lagrangian for the other cases has the same form as has been shown in [15, 18] and other papers. The precise derivation can be found in [18] and the Majorana fermion case below in Sect. 3.3.

In terms of the quantities $u_\mu, f_{\pm\mu\nu}, \chi_\pm$ defined below for each case the lowest order Lagrangian is

$$\mathcal{L}_2 = \frac{F^2}{4} \langle u_\mu u^\mu + \chi_+ \rangle. \quad (13)$$

Here we use the notation $\langle A \rangle = \text{tr}_F(A)$, denoting the trace over flavours. The NLO Lagrangian derived by [11] reads

$$\begin{aligned} \mathcal{L}_4 = & L_0 \langle u^\mu u^\nu u_\mu u_\nu \rangle + L_1 \langle u^\mu u_\mu \rangle \langle u^\nu u_\nu \rangle + L_2 \langle u^\mu u^\nu \rangle \langle u_\mu u_\nu \rangle + L_3 \langle u^\mu u_\mu u^\nu u_\nu \rangle \\ & + L_4 \langle u^\mu u_\mu \rangle \langle \chi_+ \rangle + L_5 \langle u^\mu u_\mu \chi_+ \rangle + L_6 \langle \chi_+ \rangle^2 + L_7 \langle \chi_- \rangle^2 + \frac{1}{2} L_8 \langle \chi_+^2 + \chi_-^2 \rangle \\ & - i L_9 \langle f_{+\mu\nu} u^\mu u^\nu \rangle + \frac{1}{4} L_{10} \langle f_+^2 - f_-^2 \rangle + \frac{1}{2} H_1 \langle f_+^2 - f_-^2 \rangle + \frac{1}{4} H_2 \langle \chi_+^2 - \chi_-^2 \rangle. \end{aligned} \quad (14)$$

The NNLO Lagrangian has been classified for the N_F -flavour case in [40]. The Lagrangian at NNLO for the other cases is not known, the direct equivalent of the results in [40] is definitely a complete Lagrangian but might not be minimal. For this reason we do not quote the dependence on the NNLO Lagrangian in the real and pseudo-real cases.

The divergences at NLO were derived for the QCD case in [11], for the others in [16, 18]. At NNLO only the QCD case is known [41].

3.2 The three Dirac fermion cases

A more extensive discussion can be found in [15, 18]. Here we simply quote the results.

When we have a global symmetry group G with generators T^a which is spontaneously broken down to a subgroup H with generators Q^a which form a subset of the T^a , the Goldstone bosons can be described by the coset G/H . This coset can be parametrized [42] via the broken generators X^a . Below we explain what is used for the different cases. We always work with generators normalized to 1, i.e. $\langle X^a X^b \rangle = \delta^{ab}$.

The quantities used from the quark level are given in Sect. 2.

QCD or complex representation The Goldstone boson manifold is in this case $SU(N_F) \times SU(N_F)/SU(N_F)$ which itself has the structure of an $SU(N_F)$. Note that the axial generators do not generate a subgroup of $SU(N_F) \times SU(N_F)$ even if G/H has the structure of a group in this case.

We choose as the broken generators X^a the generators of $SU(N_F) \approx G/H$. The quantities needed to construct the Lagrangian and their symmetry transformations are

$$\begin{aligned} u &= \exp\left(\frac{i}{\sqrt{2}F} \pi^a X^a\right) \rightarrow g_R u h^\dagger \equiv h u g_L^\dagger \\ u_\mu &= i(u^\dagger(\partial_\mu - i r_\mu)u - u(\partial_\mu - l_\mu)u^\dagger) \rightarrow h u_\mu h^\dagger, \\ \chi &= 2B_0 \mathcal{M} \rightarrow g_R \chi g_L^\dagger \\ \chi_\pm &= u^\dagger \chi u^\dagger \pm u \chi^\dagger u \rightarrow h \chi_\pm h^\dagger, \end{aligned}$$

$$\begin{aligned}
l_{\mu\nu} &= \partial_\mu l_\nu - \partial_\nu l_\mu - il_\mu l_\nu + il_\nu l_\mu \rightarrow g_L l_{\mu\nu} g_L^\dagger \\
r_{\mu\nu} &= \partial_\mu r_\nu - \partial_\nu r_\mu - ir_\mu r_\nu + ir_\nu r_\mu \rightarrow g_R r_{\mu\nu} g_R^\dagger \\
f_{\pm\mu\nu} &= ul_{\mu\nu}u^\dagger \pm u^\dagger r_{\mu\nu}u \rightarrow hf_{\pm\mu\nu}h^\dagger.
\end{aligned} \tag{15}$$

The first line defines h [42].

Adjoint or real representation The Goldstone boson manifold is in this case $SU(2N_F)/SO(2N_F)$. The unbroken generators satisfy $Q^a J_S = -J_S Q^{aT}$ which follows from $g J_S g^T = J_S$. The broken generators satisfy $J_S X^a = X^{aT} J_S$.

The quantities needed to construct the Lagrangians are [18]

$$\begin{aligned}
u &= \exp\left(\frac{i}{\sqrt{2}F}\pi^a X^a\right) \rightarrow guh^\dagger \\
u_\mu &= i\left(u^\dagger(\partial_\mu - i\hat{V}_\mu)u - u(\partial_\mu + iJ_S\hat{V}_\mu^T J_S)u^\dagger\right), \\
\chi &= 2B_0\hat{\mathcal{M}} \\
\chi_\pm &= u^\dagger\chi J_S u^\dagger \pm u J_S \chi^\dagger u \\
\hat{V}_{\mu\nu} &= \partial_\mu \hat{V}_\nu - \partial_\nu \hat{V}_\mu - i(\hat{V}_\mu \hat{V}_\nu - \hat{V}_\nu \hat{V}_\mu) \\
f_{\pm\mu\nu} &= J_S u \hat{V}_{\mu\nu} u^\dagger J_S \pm u \hat{V}_{\mu\nu} u^\dagger
\end{aligned} \tag{16}$$

The first line defines h by requiring that guh^\dagger is of the form $\exp(i\pi^a X^a/(\sqrt{2}F))$. Note that the derivation used $J_S u = u^T J_S$.

$N_c = 2$ or pseudo-real representation The Goldstone boson manifold is $SU(2N_F)/Sp(2N_F)$. The unbroken generators satisfy $Q^a J_A = -J_A Q^{aT}$ which follows from $g J_A g^T = J_A$. The broken generators satisfy $J_A X^a = X^{aT} J_A$.

The quantities needed are [18]

$$\begin{aligned}
u &= \exp\left(\frac{i}{\sqrt{2}F}\pi^a X^a\right) \rightarrow guh^\dagger \\
u_\mu &= i\left(u^\dagger(\partial_\mu - i\hat{V}_\mu)u - u(\partial_\mu + iJ_A\hat{V}_\mu^T J_A^T)u^\dagger\right), \\
\chi &= 2B_0\hat{\mathcal{M}} \\
\chi_\pm &= u^\dagger\chi J_A^T u^\dagger \pm u J_A \chi^\dagger u \\
\hat{V}_{\mu\nu} &= \partial_\mu \hat{V}_\nu - \partial_\nu \hat{V}_\mu - i(\hat{V}_\mu \hat{V}_\nu - \hat{V}_\nu \hat{V}_\mu) \\
f_{\pm\mu\nu} &= J_A u \hat{V}_{\mu\nu} u^\dagger J_A^T \pm u \hat{V}_{\mu\nu} u^\dagger
\end{aligned} \tag{17}$$

The first line defines h by requiring that guh^\dagger is of the form $\exp(i\pi^a X^a/(\sqrt{2}F))$. Note that the derivation used $J_A u = u^T J_A$.

3.3 Majorana fermions in a real representation

The vacuum in this case is characterized by the condensate

$$\frac{1}{2}\langle\hat{q}_i^T C\hat{q}_j\rangle = \frac{1}{2}\langle\bar{q}q\rangle\delta_{ij}. \quad (18)$$

Under the symmetry group $g \in SU(N_F)$ this moves around as

$$\delta_{ij} \rightarrow (g^T g)_{ij}. \quad (19)$$

The unbroken part of the group is given by the generators \tilde{Q}^a and the broken part by the generators \tilde{X}^a which satisfy

$$\tilde{Q}^a = -\tilde{Q}^{aT}, \quad \tilde{X}^a = \tilde{X}^{aT}. \quad (20)$$

Just as in the cases discussed in [18] we can construct a rotated vacuum in general by using the broken part of the symmetry group on the vacuum. This leads to a matrix

$$U = uu^T \rightarrow gUg^T \quad \text{with} \quad u = \exp\left(\frac{i}{\sqrt{2}F}\pi^a X^a\right). \quad (21)$$

The matrix u transforms as in the general $CCWZ$ case as

$$u \rightarrow guh^\dagger. \quad (22)$$

Some earlier work used the matrix U to describe the Lagrangian [15]. Here we will, as in [18] use the $CCWZ$ scheme to obtain a notation that is formally identical to the QCD case. We add $N_F \times N_F$ matrices of external fields \hat{V}_μ and $\hat{\mathcal{M}}$. We need to obtain the u_μ , or broken generator, parts of $u^\dagger(\partial_\mu - iV_\mu)u$. Eq. (20) have as a consequence that u satisfies

$$u = u^T. \quad (23)$$

This leads using the same method as in [18] to

$$u_\mu = i\left(u^\dagger(\partial_\mu - i\hat{V}_\mu)u - u(\partial_\mu + i\hat{V}_\mu^T)u^\dagger\right). \quad (24)$$

With this we can construct Lagrangians. The equivalent quantities to the field strengths are

$$f_{\pm\mu\nu} = u\hat{V}_{\mu\nu}u^\dagger \pm u\hat{V}_{\mu\nu}u^\dagger \quad (25)$$

with $\hat{V}_{\mu\nu} = \partial_\mu\hat{V}_\nu - \partial_\nu\hat{V}_\mu - i(\hat{V}_\mu\hat{V}_\nu - \hat{V}_\nu\hat{V}_\mu)$ and for the mass matrix

$$\chi_\pm = u^\dagger\chi u^{\dagger T} \pm u^T\chi^\dagger u \quad (26)$$

with $\chi = 2B_0\hat{\mathcal{M}}$. The Lagrangians at LO and NLO have exactly the same form as given in (13) and (14) with u_μ , χ_\pm and $f_{\pm\mu\nu}$ as defined in (24), (25) and (26).

4 Relation Dirac and Majorana for the adjoint case

As discussed below, we have calculated the adjoint case using two methods. They were appropriate for the Dirac and the Majorana case respectively. After doing the trivial $2N_F \rightarrow N_F$ change the results agreed exactly. If we compare the two cases, we see that the main difference is really the choice of vacuum.

The Dirac and Majorana cases lead to a choice of vacuum

$$\langle \hat{q}_i^T C \hat{q}_j \rangle_D \propto J_{Sij}, \quad \langle \hat{q}_i^T C \hat{q}_j \rangle_D \propto I_{ij}. \quad (27)$$

Is it possible to relate the two cases in a simple way? Under a global symmetry transformation the first one transforms as $J_S \rightarrow g J_S g^T$. If we could find a global transformation g_R that lead to $g_R J_S g_R^T = I$ the two cases would be obviously the same.

It is not possible in general with a $SU(2N_F)$ rotation to accomplish this since $\det J_S = \pm 1$ (-1 for the $2N_F = 2$) while $\det I = 1$. However it is possible with a $U(2N)$ transformation. An explicit choice for g_R , with a free phase α is

$$g_R = \frac{1}{\sqrt{2}} \begin{pmatrix} \mp i e^{i\alpha} I & \pm i e^{-i\alpha} I \\ e^{i\alpha} I & e^{-i\alpha} I \end{pmatrix}. \quad (28)$$

It can be checked that this transforms a Dirac mass term for N_F Dirac fermions into a Majorana mass term for $2N_F$ Majorana fermions.

Inspections of the effective Lagrangians needed lead to the immediate conclusion that the mass independent terms really are $U(2N_F)$ invariant, and the mass dependent terms for the two cases are turned into each other.

g_R can also be used to relate the two different embeddings of $SO(2N_F)$ in $SU(2N_F)$ to each other. For the Dirac case the $SO(2N)$ generators satisfied $Q^{aT} J_S = -J_S Q^a$ while for the Majorana case they satisfied $\tilde{Q}^{aT} = -\tilde{Q}^a$. The two sets of generators are related by

$$\tilde{Q}^a = g_R Q^a g_R^\dagger, \quad \tilde{X}^a = g_R X^a g_R^\dagger. \quad (29)$$

5 Partially quenching and the quark flow technique

A thorough discussion of PQChPT and in particular the derivation of the propagator used there is [43]. That discussion uses the supersymmetric method. Alternative methods of calculation are the replica trick [44] and the quark flow method [24]. The earliest partially quenched work for QCDlike theories used the supersymmetric method [17]. The replica trick has been used in [45]. We use the quark-flow method.

For this method we look at the matrix

$$\Phi = \pi^a X^a \quad (30)$$

for each of the cases.

For the QCD case, Φ is a traceless Hermitian matrix. We actually keep Φ in the flavour basis with elements ϕ_{ij} and i, j are flavour indices. The tracelessness condition is enforced

by the propagator. The indices are kept explicitly and the propagator connecting a field ϕ_{ij} to ϕ_{kl} is [43]

$$G_{ijkl}(k) = G_{ij}^c(k)\delta_{il}\delta_{jk} - \delta_{ij}\delta_{kl}G_{ik}^q(k)/n_{\text{sea}}. \quad (31)$$

The number of sea quarks n_{sea} is what we call N_F . with $G_{ij}^c = i/(p^2 - \chi_{ij})$. The neutral part of the propagator, G_{ik}^q , can contain double poles. In particular for the mass cases we consider:

$$\begin{aligned} G_{vv'}^q &= i(\chi_1 - \chi_4)/(p^2 - \chi_1)^2 + i/(p^2 - \chi_1), \\ G_{vs}^q &= i/(p^2 - \chi_1), \\ G_{ss'}^q &= i/(p^2 - \chi_4). \end{aligned} \quad (32)$$

v, s denote valence or sea quarks. The extra parts come from integrating out the Φ_0 [43] and enforce the condition that Φ must be traceless. When constructing the Feynman diagrams, we keep all flavour indices free. Those that connect to external states get replaced by the value of the external valence flavour index and the remaining ones are summed over the sea quark flavours. In the present calculation, with all sea quarks the same mass, that corresponds to a factor of N_F for each free flavour index.

For the Majorana, $SU(N_F) \rightarrow SO(N_F)$, case we have that $\Phi = \pi^a X^a$ with Φ Hermitian, traceless and symmetric. Hermitian and traceless follow from $SU(N_F)$ and symmetric from (20). Going to the flavour basis for the diagonal elements of Φ there is no change w.r.t. the QCD case, but the flavour charged or off-diagonal elements must be correctly symmetrized. This has to be done both for the propagator and the connection to the external states, keeping track of the needed normalization. Afterwards we set the flavour indices connected to external states to their valence values and sum over the flavours for the free indices.

For the Dirac adjoint case, $SU(2N_F) \rightarrow SO(2N_F)$, case we have that $\Phi = \pi^a X^a$ with Φ Hermitian, traceless and satisfying $X^a J_S = J_S X^{aT}$ and the matrix Φ is $2N_F \times 2N_F$. Rewriting Φ with $N_F \times N_F$ matrices leads to the form

$$\Phi = \begin{pmatrix} \Phi_A & \Phi_C^\dagger \\ \Phi_C & \Phi_A^T \end{pmatrix}, \quad \text{with } \langle A \rangle = 0, \quad \Phi_C = \Phi_C^T. \quad (33)$$

Φ_A is Hermitian. The elements in Φ_A correspond to quark-antiquark states, those in Φ_C to diquark states. Φ_A can be treated exactly as in the QCD case, both the diagonal and flavour charged or offdiagonal elements, since $\langle \Phi_A \rangle = 0$ replaces $\langle \Phi \rangle = 0$ in the QCD case. Φ_C can be treated as offdiagonal or flavour charged propagators but the needed symmetrizing should be taken care of both for external states and propagators. The normalization of all states must be done correctly as well. After constructing Feynman diagrams with both Φ_A and Φ_C degrees of freedom taken into account, we sum free index lines over the N_F degrees of freedom, not $2N_F$. The results always agree with the calculations done with the previous, Majorana, method.

For the last case, $SU(2N_F) \rightarrow Sp(2N_F)$, pseudo-real, we have that $\Phi = \pi^a X^a$ with Φ Hermitian, traceless and satisfying $X^a J_A = J_A X^{aT}$ and the matrix Φ is $2N_F \times 2N_F$.

Rewriting Φ with $N_F \times N_F$ matrices leads to the form

$$\Phi = \begin{pmatrix} \Phi_A & \Phi_C^\dagger \\ \Phi_C & \Phi_A^T \end{pmatrix}, \quad \text{with } \langle A \rangle = 0, \quad \Phi_C = -\Phi_C^T. \quad (34)$$

Φ_A is Hermitian. The elements in Φ_A correspond to quark-antiquark states, those in Φ_C to diquark states. Φ_A can be treated exactly as in the QCD case, both the diagonal and flavour charged or offdiagonal elements, since $\langle \Phi_A \rangle = 0$ replaces $\langle \Phi \rangle = 0$ in the QCD case. Φ_C can be treated as offdiagonal or flavour charged propagators but the needed antisymmetrizing should be taken care of. The normalization of all states must be done correctly as well. After constructing Feynman diagrams with both Φ_A and Φ_C degrees of freedom taken into account, we sum free index lines over the N_F degrees of freedom, not $2N_F$. In this case and the previous we can also compare calculations with Φ_A or Φ_C external states providing a check on our results.

6 Analytical results

We have calculated the masses, decay constants and vacuum expectation values to NNLO for the QCD-like theories with the symmetry breaking patterns discussed above. A number of checks have been performed on the analytical formulas. The infinite volume unquenched results were obtained earlier in [18] and we have reproduced those. The partially quenched and finite volume results in the QCD case are finite. The partially quenched expressions reduce to the unquenched results whenever we set the sea mass equal to the valence mass. In addition we reproduce the known results at NLO for the condensate [17] also for the partially quenched case. The finite volume expressions have been checked against the known NLO results and numerically with the earlier known NNLO results, as discussed in Sect. 7.

For the real and pseudo-real case we have the additional check that calculating the mass or decay constant of a quark-anti-quark or a diquark meson gives the same results. This corresponds to using a field from the A or the C sector in the matrices (33,34). For the real case we have the additional check that the results using the Dirac case and the Majorana case coincide.

The finite volume case is always done for three spatial dimensions of size L and an infinite temporal volume. In addition we work in the center of mass system, the momenta are such that the external states have zero spatial momentum.

The masses are the physical masses as defined as the pole of the full propagator. We consider here the case where all valence quarks have the same quark mass $m_1 = \hat{m}$ and the sea quarks all have the same mass $m_4 = m_S$. For the unquenched case obviously $m_4 = m_1$. The labeling is similar to those used in three flavour PQChPT [25, 26, 27, 36]. In the formulas we use instead the quantities

$$\chi_1 = 2B_0 m_1, \quad \chi_4 = 2B_0 m_4, \quad \chi_{14} = \frac{1}{2} (\chi_1 + \chi_4). \quad (35)$$

These quantities are referred to in [37] as m11, m44 and m14 respectively.

The formulas are given for the cases $SU(N_F) \times SU(N_F) \rightarrow SU(N_F)$, $SU(N_F) \rightarrow SO(N_F)$ and $SU(2N_F) \rightarrow Sp(2N_F)$. Note the difference in convention for the second case compared to [18]. The three cases are referred to in the formulas with SUN, SON and SPN for the unquenched case and PQSUN, PQSON and PQSPN for the partially quenched case. In the latter case N_F refers to the number of sea quarks.

For the mass we consider a meson made of a different quark and anti-quark or a diquark state with two different quarks. These are always valence quarks. The physical mass at finite volume is given by

$$m_{\text{phys}}^2 = \chi_1 + m^{(4)2} + \Delta^V m^{(4)2} + m^{(6)2} + \Delta^V m^{(6)2}. \quad (36)$$

The superscript (n) labels the order p^n correction and Δ^V indicates the finite volume corrections. In all cases the lowest order mass squared is given by χ_1 . A further break up is done for the LEC dependent parts via the L_i^r (NLO) and K_i^r (NNLO) and the remainder via

$$\begin{aligned} m^{(4)2} &= m^{L(4)2} + m^{R(4)2} \\ m^{(6)2} &= m^{K(6)2} + m^{L(6)2} + m^{R(6)2} \\ \Delta^V m^{(6)2} &= \Delta^V m^{L(6)2} + \Delta^V m^{R(6)2} \end{aligned} \quad (37)$$

All quantities are given explicitly in [37].

The decay constant F_{phys} for the same mesons as above is expanded w.r.t. the lowest order as

$$F_{\text{phys}} = F_{LO} \left(1 + F^{(4)} + \Delta^V F^{(4)} + F^{(6)2} + \Delta^V F^{(6)} \right), \quad (38)$$

with a similar split

$$\begin{aligned} F^{(4)} &= F^{L(4)} + F^{R(4)} \\ F^{(6)} &= F^{K(6)} + F^{L(6)} + F^{R(6)} \\ \Delta^V F^{(6)} &= \Delta^V F^{L(6)} + \Delta^V F^{R(6)} \end{aligned} \quad (39)$$

All quantities are given explicitly in [37].

The vacuum expectation value is expanded in exactly the same way

$$v_{\text{phys}} = v_{LO} \left(1 + v^{(4)} + \Delta^V v^{(4)} + v^{(6)2} + \Delta^V v^{(6)} \right), \quad (40)$$

with a similar split

$$\begin{aligned} v^{(4)} &= v^{L(4)} + v^{R(4)} \\ v^{(6)} &= v^{K(6)} + v^{L(6)} + v^{R(6)} \\ \Delta^V v^{(6)} &= \Delta^V v^{L(6)} + \Delta^V v^{R(6)} \end{aligned} \quad (41)$$

All quantities are given explicitly in [37].

The quantities with K for the SON and SPN case have been set to zero. They are polynomials up to the needed degree in χ_1 and χ_4 , with an overall factor of χ_1 for the mass.

The decay constant and the vacuum expectation value were defined implicitly in [18] using a generator X^a in the axial current normalized to one and an element in $\hat{\mathcal{M}}$ normalized to one. The consequence was that in [18] $F_{LO} = F$ and $v_{LO} = -B_0 F^2$ for all cases. This is not exactly what was done in earlier work leading to differences in factors of 2 and $\sqrt{2}$. Below we explicitly specify all definitions in terms of the quark fields.

QCD or complex representation If we label the first Dirac (valence) quark by 1 and the second by 2 the decay constant and vacuum expectation value are defined as

$$\begin{aligned}\langle 0 | \bar{q}_1 \gamma_\mu \gamma_5 q_2 | M(p) \rangle &= i\sqrt{2} F_{\text{phys}} p_\mu \\ \langle \bar{q}_1 q_1 \rangle &= \langle \bar{q}_{L1} q_{R1} + \bar{q}_{R1} q_{L1} \rangle = v_{\text{phys}}\end{aligned}\quad (42)$$

M denotes a meson of that quark content with momentum p .

The resulting lowest orders are

$$F_{LO} = F \quad v_{LO} = -B_0 F^2. \quad (43)$$

Adjoint or real representation Here we have to be careful how we define the physical decay constant. We can choose to do using generators normalized to one using Dirac Fermions or generators normalized to one using the \hat{q}_i elements.

With a Dirac fermion definition, the first Dirac (valence) quark labeled by 1 and the second by 2, the definitions are

$$\begin{aligned}\langle 0 | \bar{q}_1 \gamma_\mu \gamma_5 q_2 | M(p) \rangle &= i\sqrt{2} F_{\text{phys}} p_\mu \\ \langle \bar{q}_1 q_1 \rangle &= \langle \bar{q}_{L1} q_{R1} + \bar{q}_{R1} q_{L1} \rangle = v_{\text{phys}}\end{aligned}\quad (44)$$

M denotes a meson of that quark content with momentum p . The resulting lowest orders are

$$F_{LO} = \sqrt{2} F \quad v_{LO} = -2B_0 F^2. \quad (45)$$

If we instead choose to use the Majorana case, the natural definition of the decay constant and vacuum expectation value with the first (valence) Majorana fermion labeled as 1 and the second as 2 via

$$\begin{aligned}\frac{1}{2\sqrt{2}} \langle 0 | \hat{q}_1^* \bar{\sigma}_\mu \hat{q}_2 + \hat{q}_2^* \bar{\sigma}_\mu \hat{q}_1 | M(p) \rangle &= i\sqrt{2} F_{\text{phys}} p_\mu \\ \frac{1}{2} \langle \hat{q}_1 \sigma^2 \hat{q}_1 + \hat{q}_1^* \sigma^2 \hat{q}_1^* \rangle &= v_{\text{phys}}\end{aligned}\quad (46)$$

The resulting lowest orders are

$$F_{LO} = F \quad v_{LO} = -B_0 F^2. \quad (47)$$

$N_c = 2$ or **pseudo-real representation** Here we again need to be careful how we define the physical decay constant. We can choose to do using generators normalized to one using the original Dirac Fermions or generators normalized to one using the \hat{q}_i elements.

With a Dirac fermion definition, the first Dirac (valence) quark labeled by 1 and the second by 2, the definitions are

$$\begin{aligned}\langle 0|\bar{q}_1\gamma_\mu\gamma_5q_2|M(p)\rangle &= i\sqrt{2}F_{\text{phys}}p_\mu \\ \langle \bar{q}_1q_1\rangle &= \langle \bar{q}_{L1}q_{R1} + \bar{q}_{R1}q_{L1}\rangle = v_{\text{phys}}\end{aligned}\quad (48)$$

M denotes a meson of that quark content with momentum p . The resulting lowest orders are

$$F_{LO} = \sqrt{2}F \quad v_{LO} = -2B_0F^2. \quad (49)$$

In terms of the \hat{q}_i the definitions are

$$\begin{aligned}\langle 0|\bar{\hat{q}}_1\gamma_\mu\bar{\hat{q}}_2 + \bar{\hat{q}}_{1+N_F}\gamma_\mu\bar{\hat{q}}_{2+N_F}|M(p)\rangle &= i\sqrt{2}F_{\text{phys}}p_\mu \\ \frac{1}{2}\langle \bar{\hat{q}}_{1+N_F,a}\epsilon_{ab}C\bar{\hat{q}}_{1,b} - \bar{\hat{q}}_{1,a}\epsilon_{ab}C\bar{\hat{q}}_{1+N_F,b} - \hat{q}_{1+N_F,a}\epsilon_{ab}C\hat{q}_{1,b} + \hat{q}_{1,a}\epsilon_{ab}C\hat{q}_{1+N_F,b}\rangle &= v_{\text{phys}}.\end{aligned}\quad (50)$$

7 Numerical examples and checks

The main aim of this work is to provide the lattice work with the formulas and programs needed to do the extrapolation to zero mass. We therefore only present some representative numerical results. The numerical programs are included in the latest version of CHIRON, [38, 39].

For the numbers presented we always use $\chi_1 = 0.14^2$ GeV², if not varied explicitly, and $F = 0.0877$ GeV as well as a subtraction scale $\mu = 0.77$ GeV. The length L for the finite volume has been chosen such that $L \times 0.14$ GeV=3 or $L \approx 4.2$ fm.

The LECs at NLO we choose to be those of the recent determination of [46] with the extra LEC $L_0^r = 0$. The NNLO constants we have always put to zero.

A number of numerical checks for the QCD case have been done. The unquenched infinite volume results for three flavours agree with the three flavour results of [47, 48]. The partially quenched results for masses and decay constants at infinite volume agree with the case $d_{\text{sea}} = 1, d_{\text{val}} = 1$ of [25, 26, 27]. The unquenched results for masses and decay constants at finite volume agree with [35]. The partially quenched results for masses and decay constants at finite volume agree with the case $d_{\text{sea}} = 1, d_{\text{val}} = 1$ of [36] and finally the unquenched finite volume results for the vacuum expectation value agree with the results of [32].

In Fig. 1 we show the mass squared for the infinite volume for all cases we have considered for three values of N_F . In general, as was already noticed in [18] the corrections are larger for the larger values of N_F . The corrections are also larger for the $SU(2N_F) \rightarrow Sp(2N_F)$ case since this correspond to a twice as large number of fermions as the other cases. The partially quenched results shown in the right column are at a fixed value of χ_1 . That explains why the corrections do not vanish for $\chi_4 = 0$.

The same types of results are shown for the decay constant in Fig. 2. The corrections are somewhat larger than for the masses but the convergence is typically somewhat better. The corrections for the vacuum expectation value shown in Fig. 3 are typically larger but with again a reasonable convergence from NLO to NNLO.

We can now make similar plots for the finite volume corrections. The overall size of them is as expected. The smallest mL is about two for the left hand sides of all plots. In the unquenched case the exponential falloff with the mass is clearly visible. The partially quenched cases contain a fixed mass scale χ_1 which is why the correction is more constant there, the stays at the $mL = 3$ point for the plots. The dips are caused by the finite volume corrections going through zero. The corrections to the mass are shown in Fig. 4, the decay constant in Fig. 5 and the vacuum expectation value in Fig 6.

8 Conclusions

We have calculated in the effective field theory for the three possible symmetry breaking patterns the NNLO order finite volume and partial quenching effects to NNLO in the expansion. The results satisfy a large number of checks agreeing analytically and numerically with earlier work that our results reduce to for some cases. The analytical part of this work relied heavily on FORM [49].

The analytical results are of reasonable length but given the total number of results we have included them as FORM output in a supplementary file. They can also be downloaded from [50].

The numerical programs have been included in CHIRON [38] version 0.54 which can be downloaded from [39]. We have presented results in a number of cases with typical QCD values of the parameters. The results are of the expected sizes from earlier work in three flavour ChPT. We hope these results will be useful for lattice studies of these alternative symmetry breaking patterns.

Acknowledgements

This work is supported in part by the Swedish Research Council grants 621-2011-5080 and 621-2013-4287. JB thanks the Centro de Ciencias de Benasque Pedro Pascual, where part of this work was done, for hospitality.

References

- [1] N. Brambilla *et al.*, Eur. Phys. J. C **74** (2014) 10, 2981 [arXiv:1404.3723 [hep-ph]].
- [2] F. Sannino, plenary talk at Lattice2015.
- [3] A. Hasenfratz, plenary talk at Lattice2015.




Figure 1: The physical mass squared divided by the lowest order mass squared for the unquenched (left) as a function of χ_1 and the partially quenched case (right) as a function of χ_4 with $\chi_1 = 0.14^2$ GeV². Other input as in the text. Shown are the NLO (p^4) and NNLO ($p^4 + p^6$) results for three values of N_F . Top line: $SU(N_F) \times SU(N_F) \rightarrow SU(N_F)$. Middle line: $SU(N_F) \rightarrow SO(N_F)$. Bottom line: $SU(2N_F) \rightarrow Sp(2N_F)$.




Figure 2: The decay constant divided by the lowest order value $F_0 = F_{LO}$ for the unquenched (left) as a function of χ_1 and the partially quenched case (right) as a function of χ_4 with $\chi_1 = 0.14^2 \text{ GeV}^2$. Other input as in the text. Shown are the NLO (p^4) and NNLO ($p^4 + p^6$) results for three values of N_F . Top line: $SU(N_F) \times SU(N_F) \rightarrow SU(N_F)$. Middle line: $SU(N_F) \rightarrow SO(N_F)$. Bottom line: $SU(2N_F) \rightarrow Sp(2N_F)$.




Figure 3: The vacuum expectation value divided by the lowest order value $v_0 = v_{LO}$ for the unquenched (left) as a function of χ_1 and the partially quenched case (right) as a function of χ_4 with $\chi_1 = 0.14^2 \text{ GeV}^2$. Other input as in the text. Shown are the NLO (p^4) and NNLO ($p^4 + p^6$) results for three values of N_F . Top line: $SU(N_F) \times SU(N_F) \rightarrow SU(N_F)$. Middle line: $SU(N_F) \rightarrow SO(N_F)$. Bottom line: $SU(2N_F) \rightarrow Sp(2N_F)$.




Figure 4: The absolute value of the finite volume correction to the physical mass squared divided by the lowest order mass squared for the unquenched (left) as a function of χ_1 and the partially quenched case (right) as a function of χ_4 with $\chi_1 = 0.14^2$ GeV². Shown are the NLO (p^4) and NNLO ($p^4 + p^6$) results for three values of N_F . Top line: $SU(N_F) \times SU(N_F) \rightarrow SU(N_F)$. Middle line: $SU(N_F) \rightarrow SO(N_F)$. Bottom line: $SU(2N_F) \rightarrow Sp(2N_F)$.




Figure 5: The absolute value of the finite volume correction to the decay constant divided by the lowest order value $F_0 = F_{LO}$ for the unquenched (left) as a function of χ_1 and the partially quenched case (right) as a function of χ_4 with $\chi_1 = 0.14^2 \text{ GeV}^2$. Shown are the NLO (p^4) and NNLO ($p^4 + p^6$) results for three values of N_F . Top line: $SU(N_F) \times SU(N_F) \rightarrow SU(N_F)$. Middle line: $SU(N_F) \rightarrow SO(N_F)$. Bottom line: $SU(2N_F) \rightarrow Sp(2N_F)$.




Figure 6: The absolute value of the finite volume correction to the vacuum expectation value divided by the lowest order value $v_0 = v_{LO}$ for the unquenched (left) as a function of χ_1 and the partially quenched case (right) as a function of χ_4 with $\chi_1 = 0.14^2 \text{ GeV}^2$. Shown are the NLO (p^4) and NNLO ($p^4 + p^6$) results for three values of N_F . Top line: $SU(N_F) \times SU(N_F) \rightarrow SU(N_F)$. Middle line: $SU(N_F) \rightarrow SO(N_F)$. Bottom line: $SU(2N_F) \rightarrow Sp(2N_F)$.

- [4] J. Kuti, PoS LATTICE **2013** (2014) 004.
- [5] R. Lewis, C. Pica and F. Sannino, Phys. Rev. D **85** (2012) 014504 [arXiv:1109.3513 [hep-ph]].
 A. Hietanen, R. Lewis, C. Pica and F. Sannino, JHEP **1407** (2014) 116 [arXiv:1404.2794 [hep-lat]].
 D. Schaich *et al.* [LSD Collaboration], arXiv:1506.08791 [hep-lat].
 T. N. da Silva, E. Pallante and L. Robroek, arXiv:1506.06396 [hep-th].
 M. P. Lombardo, K. Miura, T. J. Nunes da Silva and E. Pallante, Int. J. Mod. Phys. A **29** (2014) 25, 1445007 [arXiv:1410.2036 [hep-lat]].
 T. Appelquist *et al.*, Phys. Rev. D **85** (2012) 074505 [arXiv:1201.3977 [hep-lat]].
 T. DeGrand, Y. Liu, E. T. Neil, Y. Shamir and B. Svetitsky, Phys. Rev. D **91** (2015) 114502 [arXiv:1501.05665 [hep-lat]].
- [6] J. R. Andersen *et al.*, Eur. Phys. J. Plus **126** (2011) 81 [arXiv:1104.1255 [hep-ph]].
- [7] F. Sannino, Acta Phys. Polon. B **40** (2009) 3533 [arXiv:0911.0931 [hep-ph]].
- [8] C. T. Hill and E. H. Simmons, Phys. Rept. **381** (2003) 235 [Erratum-ibid. **390** (2004) 553] [arXiv:hep-ph/0203079].
- [9] S. Weinberg, Physica A **96** (1979) 327.
- [10] J. Gasser and H. Leutwyler, Annals Phys. **158** (1984) 142.
- [11] J. Gasser and H. Leutwyler, Nucl. Phys. B **250** (1985) 465.
- [12] M. E. Peskin, Nucl. Phys. B **175** (1980) 197.
- [13] J. Preskill, Nucl. Phys. B **177**, 21 (1981).
- [14] S. Dimopoulos, Nucl. Phys. B **168** (1980) 69.
- [15] J. B. Kogut, M. A. Stephanov, D. Toublan, J. J. M. Verbaarschot and A. Zhitnitsky, Nucl. Phys. B **582** (2000) 477 [arXiv:hep-ph/0001171].
- [16] K. Splittorff, D. Toublan and J. J. M. Verbaarschot, Nucl. Phys. B **620** (2002) 290 [arXiv:hep-ph/0108040].
- [17] D. Toublan and J. J. M. Verbaarschot, Nucl. Phys. B **560** (1999) 259 [hep-th/9904199].
- [18] J. Bijnens and J. Lu, JHEP **0911** (2009) 116 [arXiv:0910.5424 [hep-ph]].
- [19] J. Bijnens and J. Lu, JHEP **1103** (2011) 028 [arXiv:1102.0172 [hep-ph]].
- [20] J. Bijnens and J. Lu, JHEP **1201** (2012) 081 [arXiv:1111.1886 [hep-ph]].

- [21] C. W. Bernard and M. F. L. Golterman, Phys. Rev. D **49** (1994) 486 [hep-lat/9306005].
- [22] C. Bernard and M. Golterman, Phys. Rev. D **88** (2013) 1, 014004 [arXiv:1304.1948 [hep-lat]].
- [23] S. R. Sharpe and N. Shoresh, Phys. Rev. D **62** (2000) 094503 [hep-lat/0006017].
- [24] S. R. Sharpe, Phys. Rev. D **46** (1992) 3146 [hep-lat/9205020].
- [25] J. Bijnens, N. Danielsson and T. A. Lahde, Phys. Rev. D **70** (2004) 111503 [hep-lat/0406017].
- [26] J. Bijnens and T. A. Lahde, Phys. Rev. D **71** (2005) 094502 [hep-lat/0501014].
- [27] J. Bijnens, N. Danielsson and T. A. Lahde, Phys. Rev. D **73** (2006) 074509 [hep-lat/0602003].
- [28] J. Gasser and H. Leutwyler, Phys. Lett. B **184** (1987) 83.
- [29] J. Gasser and H. Leutwyler, Phys. Lett. B **188** (1987) 477.
- [30] J. Gasser and H. Leutwyler, Nucl. Phys. B **307** (1988) 763.
- [31] G. Colangelo and C. Haefeli, Nucl. Phys. B **744** (2006) 14 [hep-lat/0602017].
- [32] J. Bijnens and K. Ghorbani, Phys. Lett. B **636** (2006) 51 [hep-lat/0602019].
- [33] P. H. Damgaard and H. Fukaya, JHEP **0901** (2009) 052 [arXiv:0812.2797 [hep-lat]].
- [34] J. Bijnens, E. Boström and T. A. Lähde, JHEP **1401** (2014) 019 [arXiv:1311.3531 [hep-lat]].
- [35] J. Bijnens and T. Rössler, JHEP **1501** (2015) 034 [arXiv:1411.6384 [hep-lat]].
- [36] J. Bijnens and T. Rössler, arXiv:1508.07238 [hep-lat].
- [37] See the supplementary file analyticalresults.txt. This can also be downloaded from [50].
- [38] J. Bijnens, Eur. Phys. J. C **75** (2015) 1, 27 [arXiv:1412.0887 [hep-ph]].
- [39] <http://www.thep.lu.se/~bijnens/chiron/>
- [40] J. Bijnens, G. Colangelo and G. Ecker, JHEP **9902** (1999) 020 [hep-ph/9902437].
- [41] J. Bijnens, G. Colangelo and G. Ecker, Annals Phys. **280** (2000) 100 [hep-ph/9907333].
- [42] S. R. Coleman, J. Wess and B. Zumino, Phys. Rev. **177** (1969) 2239; C. G. . Callan, S. R. Coleman, J. Wess and B. Zumino, Phys. Rev. **177** (1969) 2247.

- [43] S. R. Sharpe and N. Shoresh, *Phys. Rev. D* **64** (2001) 114510 [hep-lat/0108003].
- [44] P. H. Damgaard and K. Splittorff, *Phys. Rev. D* **62** (2000) 054509 [hep-lat/0003017].
- [45] J. Levinsen, *Phys. Rev. D* **67** (2003) 125009 [hep-th/0301008].
- [46] J. Bijnens and G. Ecker, *Ann. Rev. Nucl. Part. Sci.* **64** (2014) 149 [arXiv:1405.6488 [hep-ph]].
- [47] G. Amorós, J. Bijnens and P. Talavera, *Nucl. Phys. B* **568** (2000) 319 [hep-ph/9907264].
- [48] G. Amorós, J. Bijnens and P. Talavera, *Nucl. Phys. B* **585** (2000) 293 [Erratum-ibid. *B* **598** (2001) 665] [hep-ph/0003258].
- [49] J. A. M. Vermaseren, arXiv:math-ph/0010025.
- [50] <http://www.thep.lu.se/~bijnens/chpt/>