

Neutron-skin effect in direct photon and charged hadron production in Pb+Pb collisions at the LHC

Ilkka Helenius^{a,*}, Hannu Paukkunen^{b,c,d}, Kari J. Eskola^{b,c}

^a*Department of Astronomy and Theoretical Physics, Lund University, Sölvegatan 14A, SE-223 62 Lund, Sweden*

^b*University of Jyväskylä, Department of Physics, P.O. Box 35, FI-40014 University of Jyväskylä, Finland*

^c*Helsinki Institute of Physics, P.O. Box 64, FI-00014 University of Helsinki, Finland*

^d*Departamento de Física de Partículas and IGFAE, Universidade de Santiago de Compostela, E-15782 Galicia, Spain*

Abstract

A well-established observation in nuclear physics is that in neutron-rich spherical nuclei the distribution of neutrons extends farther than the distribution of protons. In this work, we scrutinize the influence of this so called neutron-skin effect on the centrality dependence of high- p_T direct photon and charged hadron production. We find that due to the estimated spatial dependence of the nuclear parton distribution functions, it will be demanding to unambiguously expose the neutron-skin effect with direct photons. However, when taking a ratio between the cross sections for negatively and positively charged high- p_T hadrons, even centrality-dependent nuclear-PDF effects cancel making this observable a better handle on the neutron skin. Up to 20 % effects can be expected for the most peripheral collisions.

Keywords: Heavy-ion collisions, Neutron-skin effect, Centrality dependence, Direct photons, High- p_T hadrons

1. Introduction

In ultra-relativistic heavy-ion collisions the concept of centrality plays an important role in phenomena such as the jet energy loss [1, 2, 3] or the systematics of azimuthal anisotropies [4, 5]. Experimentally, the centrality of a collision is usually defined according to the amount of energy seen in a specific part of the detector, typically at large pseudorapidities [4, 6, 7]: the more energy observed, the more central the collision. The theoretical centrality categorizations are based on Glauber models [8], in which the centrality is related to impact parameter (Optical Glauber) or to the number of nucleon-nucleon collisions (Monte-Carlo Glauber). While there is no direct, unambiguous relation between the experimental and theoretical prescriptions, it is yet generally accepted that a correspondence exists in collisions of two heavy nuclei. In nucleon-nucleus collisions, however, the same experimental procedure has led to rather unexpected results [9, 10, 11] and it is now commonly believed that such a centrality classification induces a non-trivial bias on the hard process whose centrality dependence was to be measured [12, 13, 14, 15, 16].

The Glauber models take the nuclear density distribution as an input and it is typically assumed to be identical for protons and neutrons. However, the measurements at lower energies indicate that the tail of the neutron density distribution extends farther than that of the proton density

[17, 18, 19]. While this so called neutron-skin (NS) effect [20] should not have a large importance in the centrality classification itself, it leads to a growth of the relative number of neutrons at high impact parameters and thereby influences the observables sensitive to electroweak effects in peripheral (large impact-parameter) collisions of two heavy nuclei. The impact of the NS effect to W^\pm production in Pb+Pb and p+Pb collisions at the LHC was studied in Ref. [21].

In this letter, we extend the study of Ref. [21] to direct-photon and charged-hadron production at high transverse momenta (p_T) in Pb+Pb collisions at the LHC. The goal is to study whether the NS effect has a measurable impact on these observables and to quantify at which centralities and kinematics (transverse momentum, rapidity) the effect would be most pronounced. Our hope is that, later on, the NS effect could help to calibrate the centrality classification in collisions involving heavy ions.

2. Centrality dependent hard-process cross section

Centrality classification is done here using the Optical Glauber model as in Refs. [21, 22]. For the nuclear density distribution we use the two-parameter Fermi (2pF) distribution

$$\rho^A(\mathbf{r}) = \rho_0^A / (1 + e^{(|\mathbf{r}|-d_A)/a_A}), \quad (1)$$

where d_A describes the radius of the nucleus and a_A the thickness of the nuclear surface (skin) in nucleus with a mass number A . To account for the NS effect the nuclear density is written as $\rho^A(\mathbf{r}) = \rho^{p,A}(\mathbf{r}) + \rho^{n,A}(\mathbf{r})$ where now

*Corresponding author

Email addresses: ilkka.helenius@thep.lu.se (Ilkka Helenius), hannu.t.paukkunen@jyu.fi (Hannu Paukkunen), kari.eskola@jyu.fi (Kari J. Eskola)

the parameters of the 2pF distribution are different for protons and neutrons. Here we use the parameters from Ref. [17], $d_{n,\text{Pb}} = 6.70 \pm 0.03$ fm and $a_{n,\text{Pb}} = 0.55 \pm 0.03$ fm, for neutrons and $d_{p,\text{Pb}} = 6.680$ fm and $a_{p,\text{Pb}} = 0.447$ fm for protons.¹

The hard-process cross section in an $A + B$ collision for a given centrality class \mathcal{C}_k corresponding to an impact parameter interval $b_k \leq b < b_{k+1}$ (where $b = |\mathbf{b}|$) can be calculated from

$$d\sigma_{AB}^{\text{hard}}(\mathcal{C}_k) = 2\pi \int_{b_k}^{b_{k+1}} db b \int d^2\mathbf{s} \sum_{i,j} T_A^i(\mathbf{s}_1) T_B^j(\mathbf{s}_2) d\sigma_{ij}^{\text{hard}}(A, B, \mathbf{s}_1, \mathbf{s}_2), \quad (2)$$

where the nuclear thickness functions $T_A^i(\mathbf{s})$ are obtained by integrating the density over the longitudinal (i.e. beam) direction,

$$T_A^i(\mathbf{s}) \equiv \int_{-\infty}^{\infty} dz \rho^{i,A}(\mathbf{r}), \quad (3)$$

and $\mathbf{s}_{1,2} = \mathbf{s} \pm \mathbf{b}/2$ are defined according to the Figure 20 of Ref. [22]. The indices i and j run over combinations $(i, j) = (\text{p}, \text{p}), (\text{p}, \text{n}), (\text{n}, \text{p})$ and (n, n) . The impact-parameter intervals required in Eq. (2) correspond to the fractions of the total inelastic cross section $\sigma_{AB}^{\text{inel}}(\sqrt{s})$, obtained as in Refs. [21, 22] by

$$\sigma_{AB}^{\text{inel}}(\sqrt{s}) = \int_{-\infty}^{\infty} d^2\mathbf{b} \left[1 - e^{-T_{AB}(\mathbf{b}) \sigma^{\text{inel}}(\sqrt{s})} \right], \quad (4)$$

where

$$T_{AB}(\mathbf{b}) \equiv \int_{-\infty}^{\infty} d^2\mathbf{s} \sum_{i,j} T_A^j(\mathbf{s}_1) T_B^i(\mathbf{s}_2). \quad (5)$$

We take $\sigma^{\text{inel}}(\sqrt{s} = 5 \text{ TeV}) = 70$ mb [23]. The spatial dependence of the hard-process cross section $d\sigma_{ij}^{\text{hard}}$ arises here from the spatial dependence of nPDFs,

$$d\sigma_{ij}^{\text{hard}}(A, B, \mathbf{s}_1, \mathbf{s}_2) = \sum_{k,l} f_k^{i/A}(x_1, Q^2, \mathbf{s}_1) \otimes f_l^{j/B}(x_2, Q^2, \mathbf{s}_2) \otimes d\hat{\sigma}^{kl \rightarrow \text{observable}}, \quad (6)$$

where $d\hat{\sigma}^{kl \rightarrow \text{observable}}$ are perturbative coefficient functions and k and l are parton flavour indices. The nPDFs appearing in Eq. (6) above are defined as

$$f_k^{i/A}(x, Q^2, \mathbf{s}) = r_k^{i/A}(x, Q^2, \mathbf{s}) f_k^i(x, Q^2), \quad (7)$$

where $f_k^i(x, Q^2)$ is the free nucleon PDF (here CT10NLO [24]) and $r_k^{i/A}(x, Q^2, \mathbf{s})$ the nuclear modification which depends on the transverse position of the nucleon inside the

nucleus.² Here we use EPS09s nuclear modifications from Ref. [22] in which

$$r_k^{p/A}(x, Q^2, \mathbf{s}) = 1 + \sum_{j=1}^4 c_k^j(x, Q^2) [T_A^p(\mathbf{s}) + T_A^n(\mathbf{s})]^j, \quad (8)$$

where the coefficients $c_k^j(x, Q^2)$ are obtained by analyzing the A -dependence of the EPS09 [25] nPDFs. The neutron PDFs $f_k^{n/A}(x, Q^2, \mathbf{s})$ are obtained from the proton PDFs $f_k^{p/A}(x, Q^2, \mathbf{s})$ by the isospin symmetry. By combining all, Eq. (2) factorizes into purely geometric- and purely momentum-dependent parts which can be evaluated separately thereby reducing the dimensions of the required numerical integrations. We use the INCNLO program [26, 27, 28, 29] to calculate the momentum-dependent parts at next-to-leading order in perturbative QCD.

3. Results

3.1. Direct photon production

Direct photons are produced either in the hard process or by the fragmentation of high- p_T partons from the hard process. To obtain the latter contribution we convolute the partonic spectra with the BFG (set II) parton-to-photon fragmentation functions (FFs) [30]. Since the photon coupling is stronger in the case of up-type quarks than with the down-type quarks, the production rate of direct photons is larger in p+p collisions than in n+n collisions. This leads to a lower per-nucleon rate of direct photons in heavy-ion collisions than in p+p collisions due to the presence of neutrons. This is often referred to as the isospin effect and it becomes important at large values of x where the valence quarks dominate. Furthermore, since the relative fraction of neutrons grows towards the edge of nucleus due to the NS effect, an additional suppression of direct photons in peripheral collisions is expected.

A canonical way to quantify the nuclear effects is to compute the nuclear modification factor, defined in a given centrality class \mathcal{C}_k of a Pb+Pb collision as

$$R_{\text{PbPb}}^\gamma(\mathcal{C}_k) = \frac{1}{T_{\text{PbPb}}(\mathcal{C}_k)} \frac{d\sigma_{\text{PbPb}}^\gamma(\mathcal{C}_k)}{dp_T d\eta} \bigg/ \frac{d\sigma_{\text{pp}}^\gamma}{dp_T d\eta}, \quad (9)$$

where the normalization is related to the amount of interacting nuclear matter,

$$T_{\text{PbPb}}(\mathcal{C}_k) = 2\pi \int_{b_k}^{b_{k+1}} db b \int d^2\mathbf{s} \sum_{i,j} T_{\text{Pb}}^j(\mathbf{s}_1) T_{\text{Pb}}^i(\mathbf{s}_2), \quad (10)$$

where the impact parameters b_k and b_{k+1} define the centrality class \mathcal{C}_k as in Eq. (2). For the cross-section calculations

¹In the analysis of Ref. [17], the proton density was taken as fixed when fitting the neutron parameters to the data. Therefore, the proton density has no uncertainty here.

²Currently, there is no coherent way to treat the PDF nuclear modifications within the Monte-Carlo Glauber model. This is actually why, in this work, we stick to the optical version of the Glauber model.




Figure 1: The nuclear modification factor for direct photon production in peripheral Pb+Pb collisions with $\sqrt{s_{\text{NN}}} = 5.0$ TeV at mid-rapidity for two centralities 70–80 % (left) and 90–100 % (right). The results are compared to the isospin effect (dotted) and MB result (long-dashed) and the centrality dependent results are shown with (solid) and without (short-dashed) the NS effect. The uncertainties from EPS09s nPDFs (light color band) and 2pF parametrization (dark color band around the solid lines) are calculated with the NS effect. Gray band shows the EPS09s uncertainty for the MB result in the right-hand panel.




Figure 2: The central-to-peripheral ratio for direct photon production in Pb+Pb collisions with $\sqrt{s} = 5.0$ TeV at mid-rapidity for (0–10 %)/(70–80 %) (upper left) and (0–10 %)/(90–100 %) (upper right), and at forward rapidity for (0–10 %)/(70–80 %) (lower left) and (0–10 %)/(90–100 %) (lower right). The results with (solid) and without (dashed) NS effects are shown and the uncertainties are from the EPS09s nPDFs (light color band) and from the 2pF-parametrization (dark color bands).

we have set the renormalization, factorization and fragmentation scales to photon p_T . The uncertainties related to

the scale ambiguities are not considered here in more detail since they largely cancel out in the ratio, especially at large

values of p_T relevant here [31]. The isolation criterion, often used by experiments to suppress secondary photons from hadronic decays, is not applied here since the effect to R_{PbPb}^γ is negligible at the very high values of p_T considered here.³

Figure 1 shows R_{PbPb}^γ at mid-rapidity for two centrality classes, 70–80 % and 90–100 % with and without the NS effect, compared also to the minimum-bias (0–100 %, MB) result and to the isospin effect. The uncertainties considered here are the EPS09s uncertainty (light colour band) and the one related to the uncertainty of neutrons 2pF parameters (dark colour band), obtained by evaluating R_{PbPb}^γ with the quoted parameter variations, and adding the differences to the central prediction in quadrature.

The different p_T regions are sensitive to different nPDF effects. First, comparing the MB-result to the result without nPDF effects (only isospin), at $p_T < 30$ GeV/c some suppression due to shadowing is observed which then turns into an enhancement due to anti-shadowing. At $p_T > 300$ GeV/c a suppression due to the EMC effect is observed. The spatial dependence of the nPDFs always decreases the nuclear effects towards more peripheral collisions whereas the NS effect generates additional suppression with increasing p_T . Therefore, at high values of p_T where the impact of NS is more pronounced, these two effects pull towards opposite directions thereby “softening” the aggregate centrality dependence.

As the nuclear modifications of the PDFs gradually disappear with increasing peripherality, also the uncertainty becomes smaller for more peripheral events. However, even in the 90–100 % bin the nPDF uncertainty is of the same order as the NS effect which further complicates the separation of different effects. The non-zero nPDF effects even at the most peripheral bin are due to the power series ansatz in EPS09s, see Eq. (8), which, by construction, gives zero nuclear modifications only when $b \rightarrow \infty$. The uncertainty from the neutron 2pF parametrization turns out to be rather small as the contribution of the n+n channel is inferior e.g. to the contribution of the p+p channel and thus the variations in the neutron density are not that important. Since the fraction of neutrons (and therefore the n+n channel contribution) grows towards more peripheral collisions, also the uncertainty grows accordingly.

The centrality dependence can also be studied using the central-to-peripheral ratio R_{CP} defined as

$$R_{\text{CP}} = \frac{T_{\text{PbPb}}(P)}{T_{\text{PbPb}}(C)} \frac{d\sigma_{\text{PbPb}}^\gamma(C)}{dp_T d\eta} \bigg/ \frac{d\sigma_{\text{PbPb}}^\gamma(P)}{dp_T d\eta}. \quad (11)$$

The advantage is that there is no need for a separate p+p baseline measurement and also that some uncertainties are expected to cancel out. Here, we have used the bin 0–10 % as the central result and compared it to the 70–80 % and 90–100 % bins. The results are shown in Figure 2 again with

³The valence quark-gluon channel dominates irrespectively of the isolation.

and without the NS effect. Since the peripheral bins are now in the denominator, the NS effect increases the ratio and therefore decreases the centrality dependence at high- p_T region (R_{CP} closer to unity). The nPDF originating uncertainties are now larger with the most peripheral bin red (90–100 %) because the uncertainties in the central bin (similar than the MB uncertainty in Figure 1) do not cancel here as effectively as with the less peripheral bin (70–80 %). Even though the interpretation of this observable is easier, the NS effect is still of the same order as the nPDF uncertainties.

At forward/backward rapidities (the lower panels in Figure 2) the nPDF uncertainties are smaller. This is because here the dominant contribution comes from $q+g$ initial state where the gluon is at shadowing region with only mild uncertainty (at high factorization scale), and the quark is also a well-constrained high- x valence quark. The modifications, however, are quite small and since there is an additional uncertainty due to modeling of the spatial dependence of the nPDFs, it is difficult to unambiguously study the NS effect with this observable. The most accurate centrality dependent measurement for photons in Pb+Pb comes from the ATLAS collaboration [32]. However, their most peripheral bin 40–80 % is still too central, and also the experimental uncertainties are large, to see any effects of NS.

3.2. Charged hadron production

An observable in which the nPDF effects should cancel out very efficiently but yet be sensitive to the NS effect, is the ratio between negatively and positively charged hadrons,

$$\frac{h^-}{h^+}(C_k) = \frac{d\sigma_{\text{PbPb}}^{h^-}(C_k)}{dp_T d\eta} \bigg/ \frac{d\sigma_{\text{PbPb}}^{h^+}(C_k)}{dp_T d\eta}. \quad (12)$$

Since the relative number of neutron-involving (p+n, n+p, n+n) collisions is higher in peripheral than in central collisions, the increased d-quark contribution produces less positively charged hadrons and more negatively charged hadrons during the fragmentation. We focus on computing this observable in the very high- p_T region ($p_T \gtrsim 100$ GeV/c) at the LHC, where the final state effects (suppression of hadrons) can be expected to be negligible, and where the most recent measurements indeed are consistent with a vanishing hadron-suppression effect [33]. Moreover, we would expect our calculation to be valid even at lower values of p_T (say, at $p_T \gtrsim 10$ GeV/c) where a significant suppression of the hadron production has been observed [34, 35] at all centralities. These measurements show that the particle ratios $(K^+ + K^-)/(\pi^+ + \pi^-)$ and $(p + \bar{p})/(\pi^+ + \pi^-)$ become the same in p+p and Pb+Pb collisions, which suggests a cancellation of the suppression effects in the ratio h^-/h^+ .

The cross section for hadron production is calculated by convoluting the partonic spectra with non-perturbative parton-to-hadron FFs. We consider three options, DSS [36],




Figure 3: **Upper left panel:** The ratio between positively and negatively charged hadrons in p+p (solid) and in n+n (dashed) collisions at $\sqrt{s} = 5.0$ TeV and $\eta = 0$ using DSS (green), Kretzer (blue), and AKK08 (red) FFs. **Upper right panel:** The ratio between positively and negatively charged hadrons in Pb+Pb collisions at $\sqrt{s} = 5.0$ TeV and $\eta = 0$ using DSS (solid) and Kretzer (dashed) FFs for centrality classes 0–100 % (red), 70–80 % (green), and 90–100 % (blue). The light colour bands show the uncertainty from the 2pF parametrization and the dark one the EPS09s uncertainty with DSS. For the 0–100 % (90–100 %) bin only the EPS09s (2pF) uncertainty is visible. **Lower panels:** The ratio between positively and negatively charged hadrons in Pb+Pb collisions at $\sqrt{s} = 5.0$ TeV and $\eta = 0$ in 70–80 % (green) and 90–100 % (blue) centralities normalized with 0–100 % using DSS (solid) and Kretzer (dashed) FFs. The colour bands show the nPDF uncertainties (dark) and uncertainty in the 2pF parametrization (light) with DSS FFs.

KRETZER [37] and AKK08 [38]. To better understand the variations seen using different FFs, the h^-/h^+ ratios in p+p and n+n collisions at $\sqrt{s} = 5.0$ TeV are shown in Figure 3. The first observation is that with AKK08 FFs the ratio in p+p actually turns negative at high- p_T , caused by the cross section for h^- becoming negative. This clearly unphysical result implies that the considered kinematic region is out of the validity region of AKK08. The results using DSS and KRETZER are not that different in p+p collisions but for n+n collisions almost a factor of two difference at the very highest values of p_T is observed. These differences between the FF analyses generate some further theoretical uncertainty for the considered observable. Turning this around, a measurement of h^-/h^+ in p+Pb or Pb+Pb collisions would clearly provide additional constraints for future FF analyses (modulo the possible final state effects in Pb+Pb).

The h^-/h^+ ratios in Pb+Pb collisions at $\sqrt{s_{NN}} = 5.0$ TeV for centrality classes 70–80 % and 90–100 % are

shown in the upper-right panel of Figure 3 together with the MB result with DSS and KRETZER FFs. The nPDF effects, including the centrality dependence and the uncertainties, are found to be negligible as expected. The uncertainty in the 2pF parametrization is negligible for the MB case but increases towards more peripheral collisions. The uncertainties are larger than in the case of direct photon production as the cross section for h^- now gets a large contribution from the n+n channel and thus carries more sensitivity to the parameter uncertainties in the neutron density. More importantly, the centrality dependence from the NS effect is clearly visible in this observable. However, the different FFs still yield rather different results but normalizing the ratio with the MB result, the FF dependence largely cancels out. This is demonstrated in the lower panels of Figure 3, where the ratios in 70–80 % and 90–100 % classes are normalized with the 0–100 % result for $\eta = 0$ and $|\eta| = 2$. Some FF dependence persists with $|\eta| = 2$ but it is still smaller or of the same order than the

uncertainty in 2pF parametrization. Also the nPDFs yield a few-percent uncertainty for the observable.

4. Summary and Outlook

We have studied the impact of the NS effect to direct photon and charged hadron production in Pb+Pb collisions at the LHC. In the case of photon production the NS effect has a 5–10% impact on R_{PbPb}^γ though the uncertainties in the nPDFs and their spatial dependence are of the same order or even larger than the expected effect. With R_{CP}^γ some of the nPDF uncertainties cancel out making the NS effect more transparent. Also, going to larger rapidities decreases the nPDF uncertainties, but still the smallness of the NS effect and the ambiguities due to the centrality dependence of the nPDFs makes the direct photon production a challenging observable to study the NS effect.

A more promising observable is the ratio between negatively and positively charged high- p_T hadrons. In this case, the spatial dependence of the nPDFs cancel out very efficiently and, in general, the NS effect has a more pronounced impact than in the case of direct photons. The downsides here are the sensitivity to the applied fragmentation functions and, towards smaller p_T , possible final-state modifications due to the produced strongly interacting medium. The first one can be cured by normalizing the ratio with the minimum bias result, but for a more detailed study of the latter, further modeling would be required. However, the disparity between the amount of initial-state up and down quarks should strongly correlate with the balance of produced negatively and positively charged hadrons, irrespectively of the exact way the produced hard partons hadronize.

We hope that in near future the NS effect could provide an additional handle to control the centrality classification and help to bridge the theoretical and experimental centrality definitions. As a further prospect, we plan to study the NS effect in the future high-luminosity lepton-ion colliders.

Acknowledgements

I. H. has been supported by the MCnetITN FP7 Marie Curie Initial Training Network, contract PITN-GA-2012-315877 and has received funding from the European Research Council (ERC) under the European Unions Horizon 2020 research and innovation programme (grant agreement No 668679). This research was supported by the European Research Council grant HotLHC ERC-2011-StG-279579 and by Xunta de Galicia (Conselleria de Educacion) – H. P. is part of the Strategic Unit AGRUP2015/11.

References

- [1] B. Abelev, et al. [ALICE collaboration], Measurement of charged jet suppression in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, JHEP 03 (2014) 013. [arXiv:1311.0633](#), [doi:10.1007/JHEP03\(2014\)013](#).

- [2] S. Chatrchyan, et al. [CMS collaboration], Jet momentum dependence of jet quenching in PbPb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, Phys. Lett. B712 (2012) 176–197. [arXiv:1202.5022](#), [doi:10.1016/j.physletb.2012.04.058](#).
- [3] G. Aad, et al. [ATLAS collaboration], Measurements of the Nuclear Modification Factor for Jets in Pb+Pb Collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS Detector, Phys. Rev. Lett. 114 (7) (2015) 072302. [arXiv:1411.2357](#), [doi:10.1103/PhysRevLett.114.072302](#).
- [4] G. Aad, et al. [ATLAS collaboration], Measurement of the centrality and pseudorapidity dependence of the integrated elliptic flow in lead-lead collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS detector, Eur. Phys. J. C74 (8) (2014) 2982. [arXiv:1405.3936](#), [doi:10.1140/epjc/s10052-014-2982-4](#).
- [5] S. Chatrchyan, et al. [CMS collaboration], Azimuthal anisotropy of charged particles at high transverse momenta in PbPb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, Phys. Rev. Lett. 109 (2012) 022301. [arXiv:1204.1850](#), [doi:10.1103/PhysRevLett.109.022301](#).
- [6] S. Chatrchyan, et al. [CMS collaboration], Observation and studies of jet quenching in PbPb collisions at nucleon-nucleon center-of-mass energy = 2.76 TeV, Phys. Rev. C84 (2011) 024906. [arXiv:1102.1957](#), [doi:10.1103/PhysRevC.84.024906](#).
- [7] B. Abelev, et al. [ALICE collaboration], Centrality determination of Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV with ALICE, Phys. Rev. C88 (4) (2013) 044909. [arXiv:1301.4361](#), [doi:10.1103/PhysRevC.88.044909](#).
- [8] M. L. Miller, K. Reygers, S. J. Sanders, P. Steinberg, Glauber modeling in high energy nuclear collisions, Ann.Rev.Nucl.Part.Sci. 57 (2007) 205–243. [arXiv:nucl-ex/0701025](#), [doi:10.1146/annurev.nucl.57.090506.123020](#).
- [9] S. Chatrchyan, et al. [CMS collaboration], Studies of dijet transverse momentum balance and pseudorapidity distributions in pPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, Eur. Phys. J. C74 (7) (2014) 2951. [arXiv:1401.4433](#), [doi:10.1140/epjc/s10052-014-2951-y](#).
- [10] G. Aad, et al. [ATLAS collaboration], Centrality and rapidity dependence of inclusive jet production in $\sqrt{s_{NN}} = 5.02$ TeV proton-lead collisions with the ATLAS detector, Phys. Lett. B748 (2015) 392–413. [arXiv:1412.4092](#), [doi:10.1016/j.physletb.2015.07.023](#).
- [11] A. Adare, et al. [PHENIX collaboration], Centrality-dependent modification of jet-production rates in deuteron-gold collisions at $\sqrt{s_{NN}}=200$ GeV, Phys. Rev. Lett. 116 (12) (2016) 122301. [arXiv:1509.04657](#), [doi:10.1103/PhysRevLett.116.122301](#).
- [12] G. Martinez-Garcia, Particle rapidity distribution in proton-nucleus collisions using the proton-contributor reference frame, [arXiv:1408.3108](#).
- [13] A. Bzdak, V. Skokov, S. Bathe, Centrality dependence of high energy jets in p+Pb collisions at energies available at the CERN Large Hadron Collider, Phys. Rev. C93 (4) (2016) 044901. [arXiv:1408.3156](#), [doi:10.1103/PhysRevC.93.044901](#).
- [14] M. Alvioli, B. A. Cole, L. Frankfurt, D. V. Perepelitsa, M. Strikman, Evidence for x -dependent proton color fluctuations in pA collisions at the CERN Large Hadron Collider, Phys. Rev. C93 (1) (2016) 011902. [arXiv:1409.7381](#), [doi:10.1103/PhysRevC.93.011902](#).
- [15] D. V. Perepelitsa, P. A. Steinberg, Calculation of centrality bias factors in p +A collisions based on a positive correlation of hard process yields with underlying event activity, [arXiv:1412.0976](#).
- [16] N. Armesto, D. C. Glhan, J. G. Milhano, Kinematic bias on centrality selection of jet events in pPb collisions at the LHC, Phys. Lett. B747 (2015) 441–445. [arXiv:1502.02986](#), [doi:10.1016/j.physletb.2015.06.032](#).
- [17] C. M. Tarbert, et al., Neutron skin of ^{208}Pb from Coherent Pion Photoproduction, Phys. Rev. Lett. 112 (24) (2014) 242502. [arXiv:1311.0168](#), [doi:10.1103/PhysRevLett.112.242502](#).
- [18] M. B. Tsang, et al., Constraints on the symmetry energy and neutron skins from experiments and theory, Phys. Rev. C86 (2012) 015803. [arXiv:1204.0466](#), [doi:10.1103/PhysRevC.86.015803](#).
- [19] J. Zenihiro, et al., Neutron density distributions of Pb-204, Pb-206, Pb-208 deduced via proton elastic scattering at $E_p=295$

- MeV, Phys. Rev. C82 (2010) 044611. doi:10.1103/PhysRevC.82.044611.
- [20] C. J. Horowitz, J. Piekarewicz, Neutron star structure and the neutron radius of Pb-208, Phys. Rev. Lett. 86 (2001) 5647. arXiv:astro-ph/0010227, doi:10.1103/PhysRevLett.86.5647.
- [21] H. Paukkunen, Neutron skin and centrality classification in high-energy heavy-ion collisions at the LHC, Phys. Lett. B745 (2015) 73–78. arXiv:1503.02448, doi:10.1016/j.physletb.2015.04.037.
- [22] I. Helenius, K. J. Eskola, H. Honkanen, C. A. Salgado, Impact-parameter dependent nuclear parton distribution functions: EPS09s and EKS98s and their applications in nuclear hard processes, JHEP 1207 (2012) 073. arXiv:1205.5359, doi:10.1007/JHEP07(2012)073.
- [23] G. Antchev, et al. [TOTEM collaboration], Luminosity-independent measurements of total, elastic and inelastic cross-sections at $\sqrt{s} = 7$ TeV, Europhys. Lett. 101 (2013) 21004. doi:10.1209/0295-5075/101/21004.
- [24] H.-L. Lai, M. Guzzi, J. Huston, Z. Li, P. M. Nadolsky, J. Pumplin, C. P. Yuan, New parton distributions for collider physics, Phys. Rev. D82 (2010) 074024. arXiv:1007.2241, doi:10.1103/PhysRevD.82.074024.
- [25] K. J. Eskola, H. Paukkunen, C. A. Salgado, EPS09 - a New Generation of NLO and LO Nuclear Parton Distribution Functions, JHEP 04 (2009) 065. arXiv:0902.4154, doi:10.1088/1126-6708/2009/04/065.
- [26] P. Aurenche, R. Baier, M. Fontannaz, D. Schiff, Prompt Photon Production at Large p(T) Scheme Invariant QCD Predictions and Comparison with Experiment, Nucl. Phys. B297 (1988) 661. doi:10.1016/0550-3213(88)90553-6.
- [27] F. Aversa, P. Chiappetta, M. Greco, J. P. Guillet, QCD Corrections to Parton-Parton Scattering Processes, Nucl. Phys. B327 (1989) 105. doi:10.1016/0550-3213(89)90288-5.
- [28] P. Aurenche, M. Fontannaz, J. P. Guillet, B. A. Kniehl, E. Pilon, et al., A Critical phenomenological study of inclusive photon production in hadronic collisions, Eur.Phys.J. C9 (1999) 107–119. arXiv:hep-ph/9811382, doi:10.1007/s100529900018.
- [29] P. Aurenche, M. Fontannaz, J. Guillet, B. A. Kniehl, M. Werlen, Large p(T) inclusive π^0 cross-sections and next-to-leading-order QCD predictions, Eur.Phys.J. C13 (2000) 347–355. arXiv:hep-ph/9910252, doi:10.1007/s100520000309.
- [30] L. Bourhis, M. Fontannaz, J. Guillet, Quarks and gluon fragmentation functions into photons, Eur.Phys.J. C2 (1998) 529–537. arXiv:hep-ph/9704447, doi:10.1007/s100520050158.
- [31] I. Helenius, K. J. Eskola, H. Paukkunen, Centrality dependence of inclusive prompt photon production in d+Au, Au+Au, p+Pb, and Pb+Pb collisions, JHEP 05 (2013) 030. arXiv:1302.5580, doi:10.1007/JHEP05(2013)030.
- [32] G. Aad, et al. [ATLAS collaboration], Centrality, rapidity and transverse momentum dependence of isolated prompt photon production in lead-lead collisions at $\sqrt{s_{NN}} = 2.76$ TeV measured with the ATLAS detector, Phys. Rev. C93 (3) (2016) 034914. arXiv:1506.08552, doi:10.1103/PhysRevC.93.034914.
- [33] CMS Collaboration, Measurement of the charged particle nuclear modification factor in PbPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, CMS-PAS-HIN-15-015.
- [34] B. B. Abelev, et al. [ALICE collaboration], Production of charged pions, kaons and protons at large transverse momenta in pp and PbPb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, Phys. Lett. B736 (2014) 196–207. arXiv:1401.1250, doi:10.1016/j.physletb.2014.07.011.
- [35] J. Adam, et al. [ALICE collaboration], Centrality dependence of the nuclear modification factor of charged pions, kaons, and protons in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, Phys. Rev. C93 (3) (2016) 034913. arXiv:1506.07287, doi:10.1103/PhysRevC.93.034913.
- [36] D. de Florian, R. Sassot, M. Stratmann, Global analysis of fragmentation functions for protons and charged hadrons, Phys. Rev. D76 (2007) 074033. arXiv:arXiv:0707.1506[hep-ph], doi:10.1103/PhysRevD.76.074033.
- [37] S. Kretzer, Fragmentation functions from flavor inclusive and flavor tagged e+ e- annihilations, Phys. Rev. D62 (2000) 054001. arXiv:hep-ph/0003177, doi:10.1103/PhysRevD.62.054001.
- [38] S. Albino, B. A. Kniehl, G. Kramer, AKK Update: Improvements from New Theoretical Input and Experimental Data, Nucl.Phys. B803 (2008) 42–104. arXiv:0803.2768, doi:10.1016/j.nuclphysb.2008.05.017.