

Testing the running coupling k_T -factorization formula for the inclusive gluon production

F.O. Durães¹, A.V. Giannini², V.P. Gonçalves^{3,4} and F.S. Navarra²

¹ *Curso de Física, Escola de Engenharia,
 Universidade Presbiteriana Mackenzie
 CEP 01302-907, São Paulo, Brazil*

² *Instituto de Física, Universidade de São Paulo,
 C.P. 66318, 05315-970 São Paulo, SP, Brazil*

³ *Department of Astronomy and Theoretical Physics,
 Lund University, SE-223 62 Lund, Sweden*

⁴ *High and Medium Energy Group,
 Instituto de Física e Matemática,
 Universidade Federal de Pelotas
 Caixa Postal 354, 96010-900, Pelotas, RS, Brazil.*

The inclusive gluon production at midrapidities is described in the Color Glass Condensate formalism using the k_T - factorization formula, which was derived at fixed coupling constant considering the scattering of a dilute system of partons with a dense one. Recent analysis demonstrated that this approach provides a satisfactory description of the experimental data for the inclusive hadron production in $pp/pA/AA$ collisions. However, these studies are based on the fixed coupling k_T - factorization formula, which does not take into account the running coupling corrections, which are important to set the scales present in the cross section. In this paper we consider the running coupling corrected k_T - factorization formula conjectured some years ago and investigate the impact of the running coupling corrections on the observables. In particular, the pseudorapidity distributions and charged hadrons multiplicity are calculated considering pp , dAu/pPb and $AuAu/PbPb$ collisions at RHIC and LHC energies. We compare the corrected running coupling predictions with those obtained using the original k_T - factorization assuming a fixed coupling or a prescription for the inclusion of the running of the coupling. Considering the Kharzeev - Levin - Nardi unintegrated gluon distribution and a simplified model for the nuclear geometry, we demonstrate that the distinct predictions are similar for the pseudorapidity distributions in $pp/pA/AA$ collisions and for the charged hadrons multiplicity in pp/pA collisions. On the other hand, the running coupling corrected k_T - factorization formula predicts a smoother energy dependence for $dN/d\eta$ in AA collisions.

PACS numbers:

Keywords: Particle production, Color Glass Condensate Formalism, k_T -factorization formula

I. INTRODUCTION

The understanding of inclusive hadron production in hadronic collisions is an important challenge for the theory of the strong interactions, since these processes are expected to be dominated by small transverse momentum exchange. In general, nonperturbative approaches and/or phenomenological models based on soft physics (e.g. Reggeon approach) are used to study hadron production with a satisfactory description of the experimental data. However, a shortcoming of these approaches is that they are not based on quarks and gluons and have no clear connection to the Quantum Chromodynamics (QCD). The QCD dynamics at high energies and large nuclei predicts the formation of a new state of matter, called Color Glass Condensate (For a review see Refs. [1]), characterized by the saturation scale Q_s , which is the typical momentum scale in the hadron wave function. The presence of this scale, which increases with energy and the atomic number, allows to treat hadron production on a solid theoretical basis, where perturbative methods can be applied. In the last years, the framework of the CGC approach have been used

to describe with success the experimental data for the hadron production. In high energy collisions we expect to observe the transition from a linear description of the QCD dynamics, based on the DGLAP [2] and/or BFKL [3] evolution equations, to a nonlinear description based on the Color Glass Condensate formalism [4]. The transition between these two regimes is determined by the saturation scale Q_s , which grows with the energy and atomic number. In the last years a comprehensive phenomenological analysis has been carried out to understand the HERA, RHIC and LHC data [1, 4]. Several theoretical studies have improved the CGC formalism by the inclusion of higher order corrections [5–8]. In particular, the running coupling corrections to the kernel of the Balitsky - Kovchegov (BK) equation [9] were calculated in Refs. [5, 6], with the solution being able to describe several observables at HERA, RHIC and LHC. More recently, the contributions of large single and double transverse momentum logarithms have been resummed to all orders and included in the BK equation [7, 8], with the resulting evolution equation being stable and generating a physically meaningful evolution of the dipole amplitude. In addition, the formalism of single inclusive hadron production in the framework of the hybrid approach proposed in Ref. [10] has been improved by the inclusion of next-to-leading order (NLO) corrections in Refs. [11–13], and a generalization to higher orders of the k_T - factorization formalism for inclusive gluon production was conjectured in Ref. [14]. As demonstrated in Ref. [13], the NLO corrections of the hybrid formalism bring a better agreement of the predictions with the LHC and RHIC data on forward hadron production. In contrast, the impact of the higher order corrections in the k_T - factorization formalism on observables is still unknown. This is the subject of the present paper.

The k_T - factorization formalism of gluon production in the central rapidity region (where the wave functions of both colliding particles are probed in the small- x regime) has been proposed in Ref. [15] and has been derived in the leading $\log(1/x)$ and fixed coupling approximations in Ref. [16], considering the scattering of a dilute parton system on a dense one (See also Ref. [17]). In a series of papers [18], Kharzeev, Levin and Nardi (KLN) have studied particle production at midrapidities in $pp/pA/AA$ collisions in terms of the k_T - factorization formalism. They have assumed that the main properties of hadron production, as for example the energy, rapidity and transverse momentum dependence, are determined in the initial stage of the collision by the interaction between gluons with transverse momentum of the order of the saturation scale Q_s . The presence of this scale regularizes the infrared behavior of the parton transverse momentum distributions and justifies a perturbative approach to the process. Since the basic predictions of the KLN approach have been qualitatively confirmed by RHIC and LHC data, several authors have improved the KLN formalism in order to obtain a quantitative description of these data. In particular, in Refs. [19–23], different models of the unintegrated gluon distribution and/or distinct treatments of the nuclear geometry have been considered. Although the k_T - factorization formula has been derived assuming that α_s is a constant, these different phenomenological studies have considered the running of the coupling constant and verified that it leads to an improvement of the agreement between theory and data. However, analysing in more detail these distinct predictions, we can observe that they were obtained using different choices of scale for the running coupling constant. Such uncertainty is expected in a leading order calculation, where the scales of the couplings are not known. Consequently, the inclusion of running coupling corrections in inclusive gluon production is an important step to obtain quantitative predictions with higher accuracy. In Ref. [14], the authors have calculated the running coupling corrections for the lowest - order gluon production cross section using the scale - setting prescription due to Brodsky, Lepage and Mackenzie (BLM) [24]. They found that the resulting cross section is expressed in terms of seven factors with running couplings, instead of the three present in the fixed coupling calculation. In particular, two of these running couplings run with complex - valued momentum scales, which are complex conjugates of each other, implying real production cross sections. Finally, this calculation fixes the scales of the running coupling constants appearing in the cross section. Based on these results for lowest - order gluon production, the authors have proposed a running coupling corrected k_T - factorization formula, which is expected to be valid in the same regime as the original fixed order formula. Although the proof of this formula is still an open question, it is expected that the resulting formula may still be a good approximation for the exact answer. Such expectation motivates the phenomenological analysis performed in this paper. In what follows we will compare the predictions of the running coupling k_T - factorization formula with those obtained assuming a fixed coupling constant and two different prescriptions for the inclusion of the running coupling in the leading order formula. In all calculations we will use the same model of the unintegrated gluon distribution and we will use the same prescription for hadronization. Moreover, we will consider a simplified treatment of the nuclear geometry. This procedure allows us to estimate more precisely the impact of the running coupling corrections on the k_T - factorization formula. Such aspects surely can and should be improved in a quantitative comparison of the formalism with the experimental data. However, we believe that our main conclusions will not be strongly modified.

This paper is organized as follows. In the next Section we will present a brief review of the k_T - factorization formalism and discuss the different prescriptions for the treatment of the coupling constant which will be considered in our analysis. In Section III, the KLN model of the unintegrated gluon distribution will be presented as well as the basic formulas for the calculation of the observables. Moreover, we will compute the pseudo - rapidity and energy distributions measured in hadron production in $pp/pA/AA$ collisions at RHIC and LHC energies. We will use the running coupling k_T - factorization formula and compare its predictions with those obtained assuming a fixed coupling

constant and two different prescriptions for the inclusion of the running coupling in the leading order formula. Finally, in Section IV we summarize our main conclusions.

II. INCLUSIVE GLUON PRODUCTION IN THE k_T -FACTORIZATION FORMALISM

In this Section we will discuss inclusive gluon production in the k_T -factorization formalism. Before presenting the main formulas, a comment is in order. As described in the Introduction, the cross section of this process was proposed in Ref. [15] and proven in Ref. [16] (See also Ref. [17]) considering the scattering of a dilute partonic system on a dense one at fixed coupling constant and in leading $\log(1/x)$ approximation. Consequently, its application is well justified for gluon production at midrapidity in pA collisions. On the other hand, in the case of pp and AA collisions at high energies, the gluon jet at midrapidities is produced by the interaction of two dense systems. In such cases, factorization breaking effects are expected to become significant [25], modifying the basic k_T -factorization formulas. However, the magnitude of these corrections is still subject of intense debate and its contribution in the kinematical range probed by the LHC is not well known. The fact that the k_T -factorization formalism allows us to obtain a very good description of the current data, suggests that these corrections are not large and that this formalism can be considered a reasonable approximation for the treatment of gluon production in pp and AA collisions at central rapidities. Therefore, in what follows, we will apply the k_T -factorization formalism to pp , pA and AA collisions.

Let us consider the production of a gluon with momentum k_T at rapidity y in a collision between the hadrons h_1 and h_2 , with $h_i = p$ or A . In the k_T -factorization formalism, the differential cross section for this process will be given by [16]

$$\frac{d^3\sigma}{d^2k_T dy} = \frac{2}{C_F} \frac{1}{\mathbf{k}^2} \int d^2q \alpha_s \phi_{h_1}(\mathbf{q}, y) \phi_{h_2}(\mathbf{k} - \mathbf{q}, Y - y), \quad (1)$$

where Y is the total rapidity interval of the collision, $C_F = (N_c^2 - 1)/2N_c$ and boldface variables denote transverse plane vectors, $\mathbf{k} = k_T = (k^1, k^2)$. Moreover, $\phi_{h_i}(x_i, \mathbf{q})$ denotes the so-called unintegrated gluon distribution, which represents the probability to find a gluon with momentum fraction x_i and transverse momentum \mathbf{q} in the hadron h_i . This distribution can be expressed as follows

$$\phi_{h_i}(\mathbf{k}, y) = \frac{C_F}{\alpha_s (2\pi)^3} \int d^2b d^2r e^{-i\mathbf{k}\cdot\mathbf{r}} \nabla_r^2 \mathcal{N}_{h_i}^G(\mathbf{r}, \mathbf{b}, y), \quad (2)$$

with $\mathcal{N}_{h_i}^G(\mathbf{r}, \mathbf{b}, y)$ being the dipole - hadron h_i forward scattering amplitude for a gluon dipole of transverse size \mathbf{r} and \mathbf{b} the impact parameter of the scattering. The behavior of \mathcal{N}^G at large rapidities (small - x) is directly related to the QCD dynamics at high energies. In the general case, it will be given by the solution of the JIMWLK evolution equation [26], but in the large- N_c limit it can be expressed in terms of the solution of the BK equation [9] for the quark dipole - hadron forward scattering amplitude. As the numerical solution of these equations including the impact parameter dependence is still very challenging [27–29], in the studies of gluon production using k_T - factorization formula the authors have introduced simplifying assumptions about the impact parameter dependence of the phenomenological models for the unintegrated gluon distributions (or about the quark dipole scattering amplitude), which are based on CGC physics and have their parameters constrained by experimental data [19–23].

In the k_T - factorization formula, Eq. (1), the cross section is proportional to the coupling constant α_s , which was assumed to be constant in its derivation. Moreover, α_s appears also in the unintegrated gluon distribution, Eq. (2). In the last years, running coupling corrections have been calculated for the BK-JIMWLK evolution equations and this allows us to estimate the contribution of these corrections to \mathcal{N}^G . However, it is still not clear how to determine the momentum scale in α_s in Eq. (2). This has motivated the generalization of Eqs. (1) and (2) by the inclusion of running coupling constant corrections [19–23]. In general, these studies assume that the factorized expression is preserved by these corrections and that the coupling constants in Eqs. (1) and (2) depend on different momentum scales. The choice of these scales is arbitrary and we found different choices in the literature. For example, in Ref. [19], the authors assume that $\alpha_s = \alpha_s(k_T^2)$ in Eq. (1) and $\alpha_s = \alpha_s(Q_s^2(x_i))$ in Eq. (2). On the other hand, in Ref. [21] it is assumed that $\alpha_s = \alpha_s(Q^2)$ in Eq. (1), with $Q^2 = \max\{k^2, (k - q)^2\}$, and the scale of running coupling in Eq. (2) is assumed to be equal to the transverse momentum of the gluon. A common characteristic of these approaches is that they assume the leading order running coupling

$$\alpha_s(Q^2) = \frac{12\pi}{\beta_0 \ln(Q^2/\Lambda_{QCD}^2)}; \quad \beta_0 = 33 - 2n_f \quad (3)$$

where Λ_{QCD} is a non-perturbative scale and n_f is the number of fermions. Moreover, very often a smooth freezing of the coupling at low scales is assumed. For example, in Ref. [19] the strong coupling is taken to be $\alpha_s(Q^2) = 0.5$

when $Q^2 \leq 0.8 \text{ GeV}^2$. As pointed out in Refs. [19, 21], the inclusion of running coupling corrections improves the description of the experimental data. However, as discussed in detail in Ref. [14], it is not clear that Eq. (1) will keep its factorized form after the inclusion of these corrections. In order to clarify this aspect, the authors of [14] have estimated the running coupling corrections in lowest-order gluon production cross section, finding that three factors of fixed coupling in lowest-order expression should be replaced by seven running couplings, with the new structure being called the *septumvirate* of couplings. Two scales of the couplings are complex-valued, but given the structure of the expression, the cross section is real. Moreover, the cross section is symmetric in the parton momentum scales. In Ref. [14] the authors have proposed a generalization of the lowest-order expression to higher-orders, which includes the small- x evolution. They proposed the following expression for the running coupling corrected k_T - factorization formula

$$\frac{d^3\sigma}{d^2k_T dy} = \frac{2C_F}{\pi^2} \frac{1}{k^2} \int d^2q \bar{\phi}_{h_1}(\mathbf{q}, y) \bar{\phi}_{h_2}(\mathbf{k} - \mathbf{q}, Y - y) \frac{\alpha_s(\Lambda_{\text{coll}}^2 e^{-5/3})}{\alpha_s(Q^2 e^{-5/3}) \alpha_s(Q^{*2} e^{-5/3})}, \quad (4)$$

with the unintegrated gluon distribution functions defined by

$$\bar{\phi}_{h_i}(\mathbf{k}, y) = \alpha_s \phi_{h_i}(\mathbf{k}, y) = \frac{C_F}{(2\pi)^3} \int d^2b d^2r e^{-i\mathbf{k}\cdot\mathbf{r}} \nabla_r^2 \mathcal{N}_{h_i}(\mathbf{r}, \mathbf{b}, y). \quad (5)$$

where Λ_{coll}^2 is a collinear infrared cutoff and the momentum scale Q is given by

$$\begin{aligned} \ln \frac{Q^2}{\mu_{\overline{\text{MS}}}^2} &= \frac{1}{2} \ln \frac{\mathbf{q}^2 (\mathbf{k} - \mathbf{q})^2}{\mu_{\overline{\text{MS}}}^4} - \frac{1}{4 \mathbf{q}^2 (\mathbf{k} - \mathbf{q})^2 [(\mathbf{k} - \mathbf{q})^2 - \mathbf{q}^2]^6} \left\{ \mathbf{k}^2 [(\mathbf{k} - \mathbf{q})^2 - \mathbf{q}^2]^3 \right. \\ &\times \left\{ [(\mathbf{k} - \mathbf{q})^2]^2 - (\mathbf{q}^2)^2 \right\} [(\mathbf{k}^2)^2 + ((\mathbf{k} - \mathbf{q})^2 - \mathbf{q}^2)^2] + 2 \mathbf{k}^2 [(\mathbf{q}^2)^3 - [(\mathbf{k} - \mathbf{q})^2]^3] \\ &- \mathbf{q}^2 (\mathbf{k} - \mathbf{q})^2 \left[2 (\mathbf{k}^2)^2 + 3 [(\mathbf{k} - \mathbf{q})^2 - \mathbf{q}^2]^2 - 3 \mathbf{k}^2 [(\mathbf{k} - \mathbf{q})^2 + \mathbf{q}^2] \right] \ln \left(\frac{(\mathbf{k} - \mathbf{q})^2}{\mathbf{q}^2} \right) \left. \right\} \\ &+ i [(\mathbf{k} - \mathbf{q})^2 - \mathbf{q}^2]^3 \left\{ \mathbf{k}^2 [(\mathbf{k} - \mathbf{q})^2 - \mathbf{q}^2] \left[\mathbf{k}^2 [(\mathbf{k} - \mathbf{q})^2 + \mathbf{q}^2] - (\mathbf{q}^2)^2 - [(\mathbf{k} - \mathbf{q})^2]^2 \right] \right. \\ &+ \mathbf{q}^2 (\mathbf{k} - \mathbf{q})^2 \left(\mathbf{k}^2 [(\mathbf{k} - \mathbf{q})^2 + \mathbf{q}^2] - 2 (\mathbf{k}^2)^2 - 2 [(\mathbf{k} - \mathbf{q})^2 - \mathbf{q}^2]^2 \right) \ln \left(\frac{(\mathbf{k} - \mathbf{q})^2}{\mathbf{q}^2} \right) \left. \right\} \\ &\times \sqrt{2 \mathbf{q}^2 (\mathbf{k} - \mathbf{q})^2 + 2 \mathbf{k}^2 (\mathbf{k} - \mathbf{q})^2 + 2 \mathbf{q}^2 \mathbf{k}^2 - (\mathbf{k}^2)^2 - (\mathbf{q}^2)^2 - [(\mathbf{k} - \mathbf{q})^2]^2} \left. \right\}, \quad (6) \end{aligned}$$

with $\mu_{\overline{\text{MS}}}^2$ being the renormalization scale in the $\overline{\text{MS}}$ scheme. Differently from Eq. (1), in the corrected expression all scales in the coupling constants are specified. Moreover, it has the expected behaviours for $\mathbf{q} \rightarrow 0$ and $\mathbf{q} \rightarrow \mathbf{k}$ [14]. In Ref. [14] the authors claim that Eq. (4), like Eq. (1), is valid both in the linear and non-linear regimes of the QCD dynamics. However, as emphasized there, only exact calculations can check the validity of this conjecture. We expect Eq. (4) to be a good approximation for the exact answer. This expectation motivates a phenomenological study using the running coupling corrected k_T - factorization formula.

III. RESULTS

In this section we will compare the predictions of the running coupling corrected k_T - factorization formula, given by Eq. (4) and denoted CF hereafter, with those derived using the original formula, Eq. (1). In the latter, we will calculate the inclusive gluon production cross section assuming a fixed value for the coupling constant (denoted FC), and also assuming that the couplings run according to the prescriptions proposed in Refs. [21] and [19], denoted hereafter by RC1 and RC2, respectively. In order to clarify the impact of the running coupling corrections in the k_T - factorization formula we will make the following approximations: (a) we will disregard the impact parameter dependence of the unintegrated gluon distributions and consider only minimum bias collisions assuming that $A_{eff} = 20$ (18.5) for Pb (Au); (b) we will assume the validity of the principle of Local Parton - Hadron Duality (LPDH), which implies that the form of the rapidity distribution for the hadron spectrum differs from the gluon spectrum only by a numerical factor. This introduces an effective mass m_h (it will be always equal to 0.350 GeV) which incorporates nonperturbative effects

and (c) we will use a single model of the unintegrated gluon distribution, namely the KLN model [18], which encodes the basic aspects of the nonlinear QCD dynamics, and is given by

$$\phi_{KLN}(\mathbf{k}, y) = \frac{2C_F}{3\pi^2\alpha_s}, \quad k \leq Q_s \quad (7)$$

$$= \frac{2C_F}{3\pi^2\alpha_s} \frac{Q_s^2}{k^2}, \quad k > Q_s, \quad (8)$$

where the saturation scale is given by $Q_s^2 = A_{eff}^{1/3} Q_0^2 (x_0/x)^\lambda$ with $Q_0 = 1$ GeV, $x_0 = 3 \times 10^{-4}$ and $\lambda = 0.288$ [30]. As in previous works [18, 19], we will multiply ϕ_{KLN} by a factor $(1-x)^4$ as prescribed by quark counting rules [31, 32] in order to simulate the behavior of the distribution at large x ($x \rightarrow 1$). All these approximations can and should be improved in a quantitative calculation of the observables. However, we believe that our simplified analysis can help us to get an insight on how the running coupling corrections (included in the k_T - factorization formula) change the observables. In what follows we will calculate the inclusive hadron production cross section, which is given by

$$\frac{d^3 N}{d\eta d^2 k_T} = \frac{K}{\sigma_s} h(y, k_T, m_h) \cdot \frac{d^3 \sigma}{d^2 k_T dy}, \quad (9)$$

where η is the pseudorapidity and $h(y, k_T, m_h)$ is the Jacobian for the conversion from rapidity to pseudorapidity, which is given by

$$h(y, k_T, m_h) = \sqrt{1 - \frac{m_h^2}{m_T^2 \cosh^2 y}}, \quad (10)$$

with $m_T^2 = k_T^2 + m_h^2$. The K -factor incorporates in an effective way the contribution of higher order corrections, of possible effects not included in the CGC formalism and also the uncertainty in the conversion of partons to hadrons. Moreover, σ_s is the average interaction area. As in previous works [19], we will correct the kinematics due to the presence of the mass scale m_h , replacing $k_T \rightarrow \sqrt{k_T^2 + m_h^2}$ in the definition of the Bjorken- x variable and also in the factor $\frac{1}{k^2}$ appearing in Eqs. (1) and (4). Moreover, we will choose $\alpha_s = 0.25$ in the fixed coupling calculations and $(N_c, n_f, \Lambda_{QCD}) = (3, 3, 0.240 \text{ GeV})$ in the RC1 and RC2 predictions. In the case of the corrected expression we will assume $\alpha_s (\Lambda_{coll}^2 e^{-5/3}) = 0.25$ and the value of $\mu_{\overline{\text{MS}}}$ will be fixed by requiring that $\mu_{\overline{\text{MS}}}^2 e^{5/3} = \Lambda_{QCD}^2$. Finally, the normalization factor K/σ_s will be treated as a free parameter to be fixed by the comparison with the experimental data at a given energy and/or rapidity.

In Fig. 1 we present our results for the pseudorapidity distributions obtained in pp collisions at different center-of-mass energies. The normalization of the different curves, given by the factor K/σ_s in Eq. (9), has been fixed at each energy in order to reproduce the experimental data at $\eta = 0$. The FC and RC1 predictions are very similar, differing of the other results at large η , whereas the RC2 curve exhibits the steepest rise (and fall) with the pseudorapidity. The CF formula yields a reasonable description of the pp data. We have verified that the necessary change of the normalization between $\sqrt{s} = 0.9$ TeV and 7 TeV is smaller than 1.0 % in the case of the CF predictions. On the other hand, for the other predictions, the change was larger than 19%.

In Fig. 2 we present our results for inclusive hadron production in dAu collisions ($\sqrt{s} = 0.2$ TeV) and in pPb collisions ($\sqrt{s} = 5.02$ TeV). The normalization of the dAu results was chosen so as to describe the data on the deuteron side and in the $\eta = 0$ region (within the error bars), simultaneously. On the other hand, in the pPb case, we normalize our predictions in such way that they reproduce the data at $\eta = 0$. As observed in our analysis of the pp results, the different models predict a distinct behavior with η , with the FC and RC1 curves being similar and the RC2 one predicting a steeper dependence. The CF prediction is able to describe the experimental data in a large range of pseudorapidities. The discrepancy appearing at large η in dAu collisions can be attributed to the simplified treatment of nuclear geometry used in our calculations. Concerning the change of the normalization K/σ_s necessary to fit data at different energies, we observe that the smallest change occurs for the CF predictions (≈ 17 %), while in the other predictions the change is of order of 40 %. A possible interpretation of this result is that the corrected formula captures important energy dependent higher - order contributions, since the same model of the unintegrated gluon distribution was used in all predictions.

In Fig. 3 we present our predictions for $AuAu$ collisions at $\sqrt{s} = 0.2$ TeV and $PbPb$ collisions at $\sqrt{s} = 2.76$ TeV. The normalization of the different curves has been fixed in order to reproduce the data at $\eta = 0$. The η dependence of the different curves is similar, with the CF one providing a reasonable description of the experimental data. In contrast with the pp and pPb/dAu cases, in heavy ion collisions the required change of the normalization, when going from one energy to another, is always large, even in the CF case [$\mathcal{O}(25\%)$].




FIG. 1: Pseudorapidity distributions measured in inclusive hadron production in pp collisions for different values of \sqrt{s} : (a) 0.9 TeV, (b) 2.36 TeV and (c) 7 TeV. Data are from Refs. [33–35].

Finally, let us compare the predictions of the different models for the energy dependence of charged hadron multiplicities at $\eta = 0$. We will consider pp , pA and AA collisions, with the pA and AA predictions being normalized by $\langle N_{\text{part}} \rangle$ and $2/\langle N_{\text{part}} \rangle$, respectively, where $\langle N_{\text{part}} \rangle$ is the average number of participants. We use the values of $\langle N_{\text{part}} \rangle$ given in Ref. [38, 43] for minimum bias pPb collisions and the 3% most central AA collisions. As we are interested in the energy dependence of the predictions, we will fix the normalization factors using the experimental data on $dN/d\eta$ in pp collisions at $\sqrt{s} = 0.9$ TeV, dAu collisions at $\sqrt{s} = 0.2$ TeV and $AuAu$ collisions at $\sqrt{s} = 0.13$ TeV. The predictions for higher energies will be parameter free. As can be seen in Fig. 4 the CF curve presents a slower growth with the energy in comparison with the predictions obtained using the original k_T -factorization formula. One has that using a simplified model for the unintegrated gluon distribution and a crude treatment of the nuclear geometry, the corrected k_T -factorization formula implies a satisfactory description of the pp and pA data. In particular, the CF predictions describe quite well the experimental data from pp collisions at high and low energies, in contrast to the other approaches that using the same inputs fail to reproduce data at $\sqrt{s} < 0.9$ TeV. Moreover, the corrected formula provides a prediction for the pA case that is closer to the experimental data. In contrast, the CF predictions underestimate the AA data for high energies, which could indicate that for such systems a more precise treatment of the unintegrated gluon distribution and nuclear geometry is fundamental to describe the data. Surely, such aspects should be investigated in the future.

IV. CONCLUSIONS

In the phenomenology of the CGC, the k_T -factorization formula, is one of the most important tools. It was originally derived assuming a fixed coupling constant and a collision between a dilute and a dense system. The




FIG. 2: Pseudorapidity distributions measured in inclusive hadron production in (a) dAu collisions ($\sqrt{s} = 0.2$ TeV) and (b) pPb collisions ($\sqrt{s} = 5.02$ TeV). Data are from Refs. [36–38].




FIG. 3: Pseudorapidity distributions measured in inclusive hadron production in $AuAu$ collisions at $\sqrt{s} = 0.2$ TeV (lower curves) and $PbPb$ collisions at $\sqrt{s} = 2.76$ TeV (upper curves). Data are from Refs. [39–42].

effect of running coupling corrections on the k_T - factorization formula was addressed in Ref. [14], where a corrected expression was proposed. The study of the implications of these corrections on the observables was analysed, for the first time, in this paper. Considering simplistic assumptions for the nuclear geometry and for the unintegrated gluon distribution, we have estimated the cross section for inclusive hadron production in $pp/pA/AA$ collisions and we have compared our results with the predictions derived from the original formula, from a fixed coupling and also from two different prescriptions for the scale choice in the running coupling constant. We demonstrated that the impact of these corrections on the observables is small, with the predictions of the distinct approaches for the pseudorapidity distributions and charged hadron multiplicities being similar. In particular, we verified that the predictions of the corrected formula yield a satisfactory description of the experimental data. The main difference arises in the energy dependence of the observables, with the corrected formula predicting a weaker energy dependence. Our results motivate more robust calculations considering a realistic unintegrated gluon distribution and a precise treatment of the nuclear geometry. Surely these aspects deserve to be investigated in more detail in the future. However, we believe that the exploratory study performed in this paper shed light on the basic implications of the corrected running coupling k_T - factorization formula and that the basic conclusions which emerge from this analysis will remain valid.




FIG. 4: Energy dependence of charged hadron multiplicities in the central region of rapidity $\eta = 0$. (a) pp , (b) pA and (c) AA collisions. The pA and AA predictions are normalized by N_{part} and $2/\langle N_{\text{part}} \rangle$, respectively. Data are from Refs. [33, 34, 38, 42–56].

Acknowledgments

The authors would like to thank A. Dumitru by useful discussions during the initial stage of this study and by useful comments in a first version of the manuscript. This work was partially financed by the Brazilian funding agencies CNPq, CAPES, FAPERGS and FAPESP.

-
- [1] N. Armesto *et al.*, J. Phys. G **35** (2008) 054001; C. A. Salgado *et al.*, J. Phys. G **39**, 015010 (2012); J. L. Albacete *et al.*, Int. J. Mod. Phys. E **22**, 1330007 (2013)
- [2] Yu. Dokshitzer, Sov. Phys. JETP **46**, 1649 (1977); V.N. Gribov and L. N. Lipatov, Sov. Nucl. Phys. **15**, 438 (1972); G. Altarelli and G. Parisi, Nucl. Phys. **B126**, 298 (1977).
- [3] L. N. Lipatov, Sov. J. Nucl. Phys. **23**, 338 (1976); E. A. Kuraev, L. N. Lipatov and V. S. Fadin, Sov. Phys. JETP **45**, 199 (1977); I. I. Balitsky and L. N. Lipatov, Sov. J. Nucl. Phys. **28**, 822 (1978).
- [4] F. Gelis, E. Iancu, J. Jalilian-Marian and R. Venugopalan, Ann. Rev. Nucl. Part. Sci. **60**, 463 (2010); E. Iancu and R. Venugopalan, arXiv:hep-ph/0303204; H. Weigert, Prog. Part. Nucl. Phys. **55**, 461 (2005); J. Jalilian-Marian and Y. V. Kovchegov, Prog. Part. Nucl. Phys. **56**, 104 (2006); J. L. Albacete and C. Marquet, Prog. Part. Nucl. Phys. **76**, 1 (2014).
- [5] I. Balitsky, Phys. Rev. D **75**, 014001 (2007); I. Balitsky and G. A. Chirilli, Phys. Rev. D **77**, 014019 (2008).
- [6] Y. V. Kovchegov and H. Weigert, Nucl. Phys. A **784**, 188 (2007); Nucl. Phys. A **789**, 260 (2007); Y. V. Kovchegov, J. Kuokkanen, K. Rummukainen and H. Weigert, Nucl. Phys. A **823**, 47 (2009).

- [7] E. Iancu, J. D. Madrigal, A. H. Mueller, G. Soyez and D. N. Triantafyllopoulos, Phys. Lett. B **750**, 643 (2015)
- [8] T. Lappi and H. Mantysaari, Phys. Rev. D **93**, 094004 (2016)
- [9] I. Balitsky, Nucl. Phys. **B463**, 99 (1996); Y. V. Kovchegov, Phys. Rev. D **60**, 034008 (1999); *ibid.* **61**, 074018 (2000).
- [10] A. Dumitru, A. Hayashigaki and J. Jalilian-Marian, Nucl. Phys. A **765**, 464 (2006)
- [11] G. A. Chirilli, B. W. Xiao and F. Yuan, Phys. Rev. Lett. **108**, 122301 (2012); A. M. Stasto, B. W. Xiao and D. Zaslavsky, Phys. Rev. Lett. **112**, 012302 (2014)
- [12] T. Altinoluk, N. Armesto, G. Beuf, A. Kovner and M. Lublinsky, Phys. Rev. D **91**, 094016 (2015)
- [13] K. Watanabe, B. W. Xiao, F. Yuan and D. Zaslavsky, Phys. Rev. D **92**, 034026 (2015)
- [14] W. A. Horowitz and Y. V. Kovchegov, Nucl. Phys. A **849**, 72 (2011).
- [15] L. V. Gribov, E. M. Levin and M. G. Ryskin, Phys. Rept. **100**, 1 (1983).
- [16] Y. V. Kovchegov and K. Tuchin, Phys. Rev. D **65**, 074026 (2002)
- [17] M. A. Braun, Phys. Lett. B **483**, 105 (2000).
- [18] D. Kharzeev and M. Nardi, Phys. Lett. B **507**, 121 (2001); D. Kharzeev and E. Levin, Phys. Lett. B **523**, 79 (2001); D. Kharzeev, E. Levin and M. Nardi, Phys. Rev. C **71**, 054903 (2005); Nucl. Phys. A **747**, 609 (2005)
- [19] E. Levin and A. H. Rezaeian, Phys. Rev. D **82**, 014022 (2010); Phys. Rev. D **82**, 054003 (2010); Phys. Rev. D **83**, 114001 (2011).
- [20] J. L. Albacete and A. Dumitru, arXiv:1011.5161 [hep-ph].
- [21] J. L. Albacete, A. Dumitru, H. Fujii and Y. Nara, Nucl. Phys. A **897**, 1 (2013).
- [22] A. Dumitru, D. E. Kharzeev, E. M. Levin and Y. Nara, Phys. Rev. C **85**, 044920 (2012).
- [23] P. Tribedy and R. Venugopalan, Nucl. Phys. A **850**, 136 (2011) Erratum: [Nucl. Phys. A **859**, 185 (2011)]; Phys. Lett. B **710**, 125 (2012) Erratum: [Phys. Lett. B **718**, 1154 (2013)]; B. Schenke, P. Tribedy and R. Venugopalan, Phys. Rev. C **89**, 024901 (2014).
- [24] S. J. Brodsky, G. P. Lepage and P. B. Mackenzie, Phys. Rev. D **28**, 228 (1983).
- [25] F. Gelis, T. Lappi and R. Venugopalan, Phys. Rev. D **78**, 054019 (2008); Phys. Rev. D **78**, 054020 (2008); Phys. Rev. D **79**, 094017 (2009).
- [26] J. Jalilian-Marian, A. Kovner, L. McLerran and H. Weigert, Phys. Rev. D **55**, 5414 (1997); J. Jalilian-Marian, A. Kovner and H. Weigert, Phys. Rev. D **59**, 014014 (1999), *ibid.* **59**, 014015 (1999), *ibid.* **59** 034007 (1999); A. Kovner, J. Guilherme Milhano and H. Weigert, Phys. Rev. D **62**, 114005 (2000); H. Weigert, Nucl. Phys. **A703**, 823 (2002); E. Iancu, A. Leonidov and L. McLerran, Nucl. Phys. **A692** (2001) 583; E. Ferreiro, E. Iancu, A. Leonidov and L. McLerran, Nucl. Phys. **A701**, 489 (2002).
- [27] E. Gotsman, M. Kozlov, E. Levin, U. Maor and E. Naftali, Nucl. Phys. A **742**, 55 (2004); A. Kormilitzin and E. Levin, Nucl. Phys. A **849**, 98 (2011); C. Contreras, E. Levin and I. Potashnikova, Nucl. Phys. A **948**, 1 (2016).
- [28] K. J. Golec-Biernat and A. M. Stasto, Nucl. Phys. B **668**, 345 (2003).
- [29] J. Berger and A. Stasto, Phys. Rev. D **83**, 034015 (2011); Phys. Rev. D **84**, 094022 (2011); JHEP **1301**, 001 (2013)
- [30] K. J. Golec-Biernat and M. Wusthoff, Phys. Rev. D **59**, 014017 (1998).
- [31] S. J. Brodsky and G. R. Farrar, Phys. Rev. Lett. **31**, 1153 (1973).
- [32] V. A. Matveev, R. M. Muradian and A. N. Tavkhelidze, Lett. Nuovo Cim. **7**, 719 (1973).
- [33] K. Aamodt *et al.* [ALICE Collaboration], Eur. Phys. J. C **68**, 89 (2010).
- [34] V. Khachatryan *et al.* [CMS Collaboration], JHEP **1002**, 041 (2010).
- [35] V. Khachatryan *et al.* [CMS Collaboration], Phys. Rev. Lett. **105**, 022002 (2010).
- [36] I. Arsene *et al.* [BRAHMS Collaboration], Phys. Rev. Lett. **94**, 032301 (2005).
- [37] B. Alver *et al.* [PHOBOS Collaboration], Phys. Rev. C **83**, 024913 (2011).
- [38] B. Abelev *et al.* [ALICE Collaboration], Phys. Rev. Lett. **110**, no. 3, 032301 (2013).
- [39] B. B. Back *et al.* [PHOBOS Collaboration], Phys. Rev. Lett. **87**, 102303 (2001).
- [40] B. B. Back *et al.*, Phys. Rev. Lett. **91**, 052303 (2003).
- [41] E. Abbas *et al.* [ALICE Collaboration], Phys. Lett. B **726**, 610 (2013).
- [42] S. Chatrchyan *et al.* [CMS Collaboration], JHEP **1108**, 141 (2011).
- [43] B. B. Back *et al.* [PHOBOS Collaboration], Phys. Rev. C **65**, 061901 (2002).
- [44] J. Whitmore, Phys. Rept. **10**, 273 (1974).
- [45] W. Thome *et al.* [Aachen-CERN-Heidelberg-Munich Collaboration], Nucl. Phys. B **129**, 365 (1977).
- [46] G. J. Alner *et al.* [UA5 Collaboration], Z. Phys. C **33**, 1 (1986).
- [47] R. Nouicer *et al.* [PHOBOS Collaboration], J. Phys. G **30**, S1133 (2004).
- [48] K. Aamodt *et al.* [ALICE Collaboration], Eur. Phys. J. C **68**, 345 (2010) doi:10.1140/epjc/s10052-010-1350-2 [arXiv:1004.3514 [hep-ex]].
- [49] C. Albajar *et al.* [UA1 Collaboration], Nucl. Phys. B **335**, 261 (1990).
- [50] B. I. Abelev *et al.* [STAR Collaboration], Phys. Rev. C **79**, 034909 (2009).
- [51] F. Abe *et al.* [CDF Collaboration], Phys. Rev. D **41**, 2330 (1990).
- [52] I. G. Bearden *et al.* [BRAHMS Collaboration], Phys. Lett. B **523**, 227 (2001).
- [53] I. G. Bearden *et al.* [BRAHMS Collaboration], Phys. Rev. Lett. **88**, 202301 (2002).
- [54] S. S. Adler *et al.* [PHENIX Collaboration], Phys. Rev. C **71**, 034908 (2005) Erratum: [Phys. Rev. C **71**, 049901 (2005)].
- [55] K. Aamodt *et al.* [ALICE Collaboration], Phys. Rev. Lett. **106**, 032301 (2011).
- [56] G. Aad *et al.* [ATLAS Collaboration], Phys. Lett. B **710**, 363 (2012).